
Adaptive Dynamics: Basic Theory and an Application

J.A.J. (= Hans) Metz (Evolution and Ecology Program, IIASA;

Institute of Biology and Department of Mathematics, Leiden University & Naturalis Biodiversity Centre, Leiden, Netherlands, Modélisation Mathématique et Biodiversité, Velolia-École Polytechnique, Paris)

An ecology-based theory of meso-evolution with practical applicability

The theory of structured populations is a mathematical framework for developing and analyzing ecological models that can take account of relatively realistic detail at the level of individual organisms. This framework has given rise to the theory of adaptive dynamics (AD), a versatile framework for dealing with the evolution of the adaptable traits of individuals subject to ecologically driven selection. By concentrating on general features of ecologies AD can be used to make general predictions about meso-evolutionary patterns. Moreover, AD has produced a general toolbox for dealing efficiently with eco-evolutionary processes of any level of complexity.




AS


* For small mutational steps Y takes over, except near so-called "ess"es.

Evolution proceeds by repeated mutant substitutions


Let the environment be defined as anything that influences the population dynamical behavior of individuals. Populations can then be represented as frequency distributions over the possible states of individuals, the movement of which depends on the environment. In a given trend-free environment populations grow on average exponentially. Fitness is the rate constant ρ of this growth. Real populations influence their environment and as a result stay bounded. Hence their dynamics moves towards an attractor which produces a trend-free environment E_{attr}. If we characterize different sorts of individuals with a trait vector, X for the resident populations and Y for potential mutants, then whether a Y mutant can invade depends on its fitness in the environment engendered by the traits X_1, \dots, X_k of the resident populations.


Some graphical tools, applicable to one-dimensional trait spaces






Applying AD ideas to the evolution of respiratory diseases





More deeply seated respiratory diseases are both less infective and more harmful. They should therefore evolve in the direction of the nose. Close to the tip they are hampered by lack of space,



making for an "optimum depth". There limited cross-immunity can lead to evolutionary branching.

Some predictions

- 1. The variety of diseases of the upper airways circulating in a population will usually be far larger than the variety of those of the lower airways.
- Since the targets of a respiratory disease will start with evolving upwards, and diversification in the vertical direction is slow, there 2. will in general be room at the bottom. Therefore, emerging respiratory diseases will usually have a deep target, be highly virulent, but will not be overly infective.