

ACCEPTED MANUSCRIPT • OPEN ACCESS

Crop productivity changes in 1.5°C and 2°C worlds under climate sensitivity uncertainty

To cite this article before publication: Carl-Friedrich Schleussner *et al* 2018 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/aab63b>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2018 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Crop productivity changes in 1.5°C and 2°C worlds under climate sensitivity uncertainty

Carl-Friedrich Schleussner^{1,2}, Delphine Deryng^{1,3}, Christoph Müller², Joshua Elliott^{3,4}, Fahad Saeed¹, Christian Folberth⁵, Wenfeng Liu⁶, Xuhui Wang^{7,8}, Thomas A. M. Pugh^{9,10}, Wim Thiery^{11,12}, Sonia I. Seneviratne¹, Joeri Rogelj⁵

¹ Climate Analytics, 10969 Berlin, Germany

² Potsdam Institute for Climate Impact Research, 14473 Potsdam, Germany

³ Columbia University Center for Climate Systems Research, New York, NY 10025, USA

⁴ University of Chicago and ANL Computation Institute, Chicago, IL 60637, USA

⁵ International Institute for Applied Systems Analysis, Ecosystem Services and Management Program, 2361 Laxenburg, Austria

⁶ Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf, Switzerland

⁷ Laboratoire des Sciences du Climat et de l'Environnement. CEA CNRS UVSQ Orme des Merisiers

⁸ Sino-French Institute of Earth System Sciences, Peking University, 100871 Beijing, China

⁹ School of Geography, Earth & Environmental Sciences and Birmingham Institute of Forest Research, University of Birmingham, B15 2TT, UK

¹⁰ Karlsruhe Institute of Technology, IMK-IFU, 82467 Garmisch-Partenkirchen, Germany

¹¹ ETH Zurich, 8092 Zürich, Switzerland

¹² Vrije Universiteit Brussel, 1050 Brussels, Belgium

Abstract

Following the adoption of the Paris Agreement, there has been an increasing interest in quantifying impacts at discrete levels of global mean temperature (GMT) increase such as 1.5°C and 2°C above pre-industrial levels. Consequences of anthropogenic greenhouse gas emissions on agricultural productivity have direct and immediate relevance for human societies. Future crop yields will be affected by anthropogenic climate change as well as direct effects of emissions such as CO₂ fertilization. At the same time, the climate sensitivity to future emissions is uncertain. Here we investigate the sensitivity of future crop yield projections with a set of global gridded crop models for four major staple crops at 1.5°C and 2°C warming above pre-industrial levels, as well as at different CO₂ levels determined by similar probabilities to lead to 1.5°C and 2°C, using climate forcing data from the Half a degree Additional warming, Prognosis and Projected Impacts (HAPPI) project. For the same CO₂ forcing, we find consistent negative effects of half a degree warming on productivity in most world regions. Increasing CO₂ concentrations consistent with these warming levels have potentially much stronger but highly uncertain effects than 0.5°C warming increments. Half a degree warming will also lead to more extreme low yields, in particular over tropical regions. Our results indicate that GMT change alone is insufficient to determine future impacts on crop productivity.

Introduction

Among the manifold impacts of anthropogenic climate change, its potential to threaten global food production has always been of particular concern (UNFCCC 1992). Observational evidence already indicates adverse impacts of climate change on crop productivity across the globe (Schlenker and Lobell 2010, Lobell *et al* 2011b, Moore and Lobell 2015) and underscore the risk posed by extreme weather events, in particular droughts and heat waves, on crop yield (Lesk *et al* 2016, Schauburger *et al* 2017, Ray *et al* 2015). In addition to changes in climatic conditions, anthropogenic greenhouse gas emissions and associated rising atmospheric CO₂ concentrations could also play a direct role on crop growth and crop yield (Kimball 2016), also related to enhanced water use efficiency (Morgan *et al* 2011). Yet CO₂ effects on crop performance are manifold as well as regionally different (McGrath and Lobell 2013, Deryng *et al* 2016), and remain a large source of uncertainty in climate impact assessment on agriculture (Asseng *et al* 2013, Rosenzweig *et al* 2014, Deryng *et al* 2016). Thus, despite the possible benefits of elevated CO₂ on crop yield, there is an emerging consensus that

adopting a stringent mitigation pathway would reduce the risks of crop yield losses, and would especially benefit agriculture and food security in the tropics and sub-tropics (Müller *et al* 2015), which face a higher risk of heat-stress damage (Lobell *et al* 2011a, Deryng *et al* 2014).

Table 1 | Applied CO₂ concentrations for the three model periods and corresponding classifications in terms of exceedance probability for the respective warming levels according to a TCR-based estimate (see Methods). Medium values correspond to standard HAPPI CO₂ concentrations.

	CO ₂ concentrations associated with different climate responses		
	Low	Medium	High
2006-2015	Observed [~390ppm]		
1.5°C	390ppm	423.4ppm	486.6ppm
2°C	423.4ppm	486.6ppm	590ppm

The adoption of the Paris Agreement and the subsequent special report of the Intergovernmental Panel on Climate Change (IPCC) on 1.5°C has led to an increasing interest in differentiation between impacts of climate change at 1.5°C above pre-industrial levels in particular in comparison to 2.0°C (Schleussner *et al* 2016b). This focus on impacts at specific warming levels calls for targeted modelling efforts (James *et al* 2017).

It also raises questions for which impacts of climate change a global mean temperature (GMT) level alone is sufficient to characterise impacts of climate change (Schleussner *et al* 2016b). In concentration scenarios such as the Representative Concentration Pathways (RCPs), CO₂ concentrations are prescribed. The climate sensitivity, however, is uncertain and differs substantially between climate models thereby leading to model-dependent warming trajectories (Stocker *et al* 2013). To account for uncertainty in the climate sensitivity, the link between CO₂ concentration pathways and GMT levels is generally explored in a probabilistic fashion (IPCC 2014). The probability for not exceeding 2°C above pre-industrial levels in the lowest RCP2.6 scenario, for example, has been assessed to be more than 66% (IPCC 2014). In a concentration pathway approach, uncertainty in the climate sensitivity is thereby consistently dealt with. For GMT focussed studies, however, the corresponding CO₂ concentrations uncertainty range has to be explored systematically. This has profound consequences for the assessment of future crop yields at specific warming levels, and the biosphere response more generally, as it is responsive both to changes in CO₂ levels as well as climate.

In the following, we assess changes in crop productivity under 1.5°C and 2°C warmer climates provided by the model intercomparison project “Half a degree Additional warming, Prognosis and Projected Impacts” (HAPPI, Mitchell *et al* 2017). Our analysis is based on modelled crop yield data from six models of the Global Gridded Crop Model Intercomparison (GGCMI, Elliott *et al* 2015, Müller *et al* 2017) as part of the Agricultural Model Intercomparison and Improvement Project (AgMIP, Rosenzweig *et al* 2013). We provide projections for the four major staple crops: wheat (*Triticum aestivum* L.), maize (*Zea mays* L.), rice (*Oryza sativa* L.), and soybean (*Glycine max* L.). Crop yield responses for varying CO₂ concentrations are analysed, which allows for disentangling the effect of CO₂ fertilization and 0.5°C warming increments. Finally, we also assess changes in 10-year minimum productivity to understand implications for yield stability - a central aspect for food security.

Methods

The HAPPI modelling protocol includes three 10-year periods with prescribed atmospheric forcing as well as sea-surface temperatures and sea-ice forcing conditions (see Mitchell *et al* 2017 for further details on the HAPPI protocol). Participating general circulation models (GCMs) have provided multi-member realisations for each of the three periods. The reference period for the HAPPI experiment is the ‘current decade’ from 2006-2015 forced by observations including observed CO₂ concentrations that have increased from 380.9 parts per million (ppm) to 402.9 ppm over this decade. Mean warming over this period corresponds to about 0.9°C above the 1860-1880 period in the Berkeley Earth GMT dataset. The Future 1.5°C experiment is based on the RCP2.6 experiment and takes constant forcing for greenhouse gases and aerosols and sea-surface temperatures from the 2091-2100 decade. CO₂ concentrations in this experiment are constant at 423.4 ppm. The Future 2°C experiment uses scaled




Figure 1: Changes in global crop productivity under 1.5°C warming (upper panel) and 2°C (lower panel) for four major staple crops (wheat, maize, soybean and rice from left to right, note that y-axis scaling is different). Projections for 7 crop models from the global gridded crop model intercomparison (GGCMI) project are shown for a set of warming levels specific CO₂ concentrations (see Table 1). The levels of CO₂ concentrations for Low, Medium and High (1.5°C: 390 ppm, 423ppm, 486ppm; 2°C: 423ppm, 486ppm, 590ppm) are chosen so that they resemble similar climate response probability levels for 1.5°C and 2°C (see Methods). Changes are derived relative to the 2006-2015 median for each GGCMI-GCM combination before aggregation. Boxes depict the interquartile range across climate-crop model multi-realisation ensembles and years (see Table S2, n=135 - 200), whiskers extend to at most 1.5 of the interquartile range. Outliers are not shown.

atmospheric and sea-surface temperature forcing from RCP2.6 and RCP4.5 with CO₂ concentrations set to 486.6 ppm.

Multi-ensemble projections for four GCMs from the HAPPI intercomparison projected have been re-gridded to a 0.5x0.5°C regular grid and bias corrected based on the EWEMBI dataset (Lange 2017) following the modelling protocol of the Intersectoral Impact Model Intercomparison Project (ISIMIP; Frieler *et al* 2016). Five bias-corrected ensemble members per GCM are used in this analysis. Harmonised agricultural management data for fertiliser application rates, irrigated and rainfed areas and crop calendar are applied according to the fully harmonized configuration (*fullharm*) as introduced in (Elliott *et al* 2015). An overview of GGCMI model setups is provided in Table S1; an overview of available GCM simulations, model years and ensemble members in Table S2. Crop producing regions are masked using rainfed and irrigated areas from the MIRCA 2000 dataset (Portmann *et al* 2010) that is also used for aggregation of crop yield over actual harvested areas (Porwollik *et al* 2017).

In addition to the core set of HAPPI experiments, the sensitivity to different CO₂ levels linked to uncertainty in the climate sensitivity is explored. A useful metric to assess the climate sensitivity to increase in CO₂ concentrations is the ‘transient climate response’ (TCR) that is defined as the annual mean GMT change at the time of CO₂ doubling following a linear increase in CO₂ forcing over a period of 70 years (Stocker *et al* 2013). The AR5 provides an estimate for a likely range for the TCR between 1°C to 2.5°C (Stocker *et al* 2013). Here we are approximating probabilities for end of century warming by this TCR estimate assuming a normal distribution with mean at 1.75°C and a standard deviation of 0.75°C. Based on this distribution, TCR probability levels for not exceeding 1.5°C and 2°C at different CO₂ concentrations are derived (see Figure S1). Radiative forcing from non-CO₂ greenhouse gases and aerosols are based on RCP2.6 (1.5°C, 0.45 W/m²) or scaled RCP2.6 and RCP4.5 (2°C, 0.63 W/m²) end of century, respectively (Mitchell *et al* 2017).

Following this TCR-based approach, the 1.5°C, non-exceedance probabilities for 390.0 ppm, 423.4 ppm and 483.0 ppm are 84%, 67% and 44%, respectively. Probabilities for 2°C and 423.4 ppm (87%) and 483.0 ppm (67%) yield quite consistent values, thereby allowing for comparing consistent GMT – CO₂ combinations. For the 2°C experiments an additional CO₂ concentration of 590.0 ppm (42%) is chosen that is in line with the high ppm-probability of the 1.5°C set. These GMT-CO₂ combinations thereby establish a consistent scenario set that in the following will be called ‘low’, ‘medium’ and ‘high’ following the respective CO₂ concentrations (see Table 1).




Figure 2: Projected changes in global (blue) and tropical (yellow) crop productivity relative to the 2006-2015 period for $\sim 0.5^\circ\text{C}$ GMT increases at different levels of CO_2 concentrations (top panel) as well as for the same warming but different CO_2 concentrations (middle and bottom panel). Top panel: For 390ppm, the difference is derived based on the 1.5°C and the 2006-2015 periods (i.e. an GMT increase of 0.7°C), for 423ppm and 486ppm between 2°C and 1.5°C . Changes in crop productivity are aggregated over present day crop producing regions following MIRCA-2000; tropics only consider these areas between 23.5°S and 23.5°N . The box-whiskers comprise the spread of mean changes for the individual GCM-GGCMs pairs (temporal and multi-realisation ensemble mean, $n=9-21$, see Table S2 for an overview of the GCM-GGCM simulations). Boxes depict the interquartile range. Note that the number of GGCMs differs per crop. Whiskers extend to at most 1.5 of the interquartile range. Outliers are not shown.

Results

The choice of CO_2 mixing ratio sets with very similar climate sensitivity probabilities for the 1.5°C and 2°C simulations allow for directly assessing the effects of climate sensitivity uncertainty on global crop productivity. Results for wheat, maize, soybean and rice are depicted in Figure 1. The response in globally aggregated crop productivity to changing CO_2 concentrations is found to be strongly model and crop dependent. For maize, which is least responsive to elevated CO_2 concentrations, most models do not indicate a substantial effect of different CO_2 levels. On the contrary, for rice as well as wheat for some models, the CO_2 level largely determines the sign of the warming effect. GGCM projections for rice indicate a change in direction of the warming impact from negative at low CO_2 level to (moderately) positive under (medium) high CO_2 levels. This dominant CO_2 effect is independent of the warming level. Results for soybean follow a similar pattern. Projections for wheat indicate generally beneficial effects of rising CO_2 levels and typically a moderately positive response to rising temperatures in most models even at the lowest CO_2 levels considered (compare Figure 1, panel 1.5°C - Low CO_2).

Regrouping of the combined GMT- CO_2 sensitivity runs allows for directly assessing the effect of $\sim 0.5^\circ\text{C}$ warming increments at different CO_2 concentrations. For the GCM ensemble used, the warming difference between the recent past (2006-2015) and the 1.5°C period is about 0.67°C , between the 1.5°C and 2°C periods around 0.45°C (see Table S3 for the GCM specific warming differences). From our set of GMT- CO_2 experiments we can thereby form three pairs to investigate the impact of $\sim 0.5^\circ\text{C}$ warming increments: 1.5°C minus recent past at 390 ppm, 2°C minus 1.5°C at 423 ppm and 2°C minus 1.5°C at 486 ppm. The resulting global as well as tropical (between $23.5^\circ\text{S}/^\circ\text{N}$) crop productivity changes are displayed in Figure 2 (top panel). Apart from a slight positive response of global productivity up to 1.5°C warming for wheat and maize, median global crop productivity is consistently negatively affected by 0.5°C warming increments. For rice, each

1
2
3 additional warming increment becomes subsequently less impactful, which is consistent with the observation
4 that simulated rice productivity changes are very responsive to elevated CO₂ concentrations (Wang *et al* 2015).
5 Differences between global and tropical yields are particularly pronounced for wheat, whereas median crop
6 productivity is projected to decrease by 2.5% as a result of additional 0.5°C warming (see also Table S4). As
7 shown in Figure 2 (middle and bottom panel), the effect of uncertainty in climate sensitivity is comparable and
8 for wheat, soy and rice more pronounced than the effect of a 0.5°C temperature increase.
9

10 Changes in crop productivity are further regionalised using the climatological regions from the IPCC SREX
11 report (IPCC 2012). Figure 3 depicts the regionally resolved changes (see also Table S4). While some high
12 latitude regions like North Asia or Northern Europe see some benefits under future warming up to 1.5°C (blue
13 bars in Figure 3), warming benefits beyond 1.5°C remain limited. Tropical and sub-tropical regions are affected
14 most strongly, with median reductions in total crop productivity of 3-5% projected for regions such as Central
15 America and the Caribbean, the Sahel or East Africa. Rice productivity is particularly affected in water-scarce
16 regions such as the Mediterranean or West Asia (projected median productivity reductions of about 5%). Future
17 drought during the summer is projected to increase (Greve and Seneviratne 2015) for these regions, the period
18 where irrigation demand is highest (Thiery *et al* 2017). This renders the projected changes for these regions
19 conservative, as no water limitations are considered for irrigated crops in our simulations. Finally, the multi-
20 ensemble nature of the HAPPI modelling protocol also allows for assessing changes in the 1-in-10 year low
21 global crop productivity as shown in Figure 4. The changes in 10yr extreme lows follow the trend for the median
22 projections displayed in Figure 2. For rice, the impact of warming from current GMT to 1.5°C is more
23 pronounced for the 1-in-10 year low harvests than the warming from 1.5 to 2.0°C at any CO₂ level.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3: As Figure 2, but aggregated over the SREX world regions. Projections are only given for regions that include at least 0.1% of global production in MIRCA-2000. Results for individual regions are also given in Table S4. Note that agricultural areas for the different regions vary substantially (see Table S5 for the regional share of grid cells with agricultural activity per crop and region).

Accepted Manuscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4: As Figure 2 top panel, but for extreme low yields with a return period of 10 years or less. Extreme low yields are derived as mean over the <10% quantile per warming level and GCM-GGCM pair (45-50 years, see Table S2). Relative changes are derived relative to the extreme low yields of the reference period.

Discussion

Our findings of consistently reducing yields under scenarios of increased warming align well with existing literature estimating the impacts of warming on crop productivity using either process-based (Rosenzweig *et al* 2014, Zhao *et al* 2017, Liu *et al* 2016) or statistical estimates (Lobell and Asseng 2017). Not considering uncertain effects of CO₂ fertilisation, median changes in local yields over the tropical crop producing regions for the four major staple crops wheat, maize, soybean and rice have been found to be negatively affected under a 1.5°C GMT increase relative to pre-industrial levels and even more so under 2°C (Schleussner *et al* 2016a). Even when accounting for the full effects of CO₂ fertilisation in crop models, median local tropical yields for wheat and maize are still found to be negatively affected and reductions to double between 1.5°C and 2°C. Our findings confirm the assessment of increasing risk for local crop productivity between 1.5°C and 2°C based on 20-year time slices at mean warming levels of 1.5°C and 2°C from RCP8.5 simulations from the ISIMIP Fast Track experiment (Warszawski *et al* 2013). If at all, our reported reductions are on the low end. For wheat, for example, we find a reduction for global productivity of about 2% per 0.5°C warming (likely range -2.7 to +0.%) compared to 4-6% per degree of warming reported in other studies combining observational and model evidence (Asseng *et al* 2014, Liu *et al* 2016). Zhao *et al* (2017) have investigated impacts of GMT increase for all four major staple crops at 380ppm. They find warming to reduce global yields of wheat by $6.0 \pm 2.9\%$, rice by $3.2 \pm 3.7\%$, maize by $7.4 \pm 4.5\%$ and soybean by $3.1 \pm 5\%$ per °C GMT increase. Our findings for soybean and rice are well within the confidence range of Zhao *et al* 2017, but our median projections for maize are again slightly more conservative.

One possible origin for our lower estimate is the limited capability of most models in our ensemble to represent the effects of heat stress on wheat that is found to play a dominant role in productivity losses in field studies (Asseng *et al* 2014) and observations (Liu *et al* 2016) and different temperature response mechanisms in models are a major source of uncertainty in wheat (Wang *et al* 2017). Similar effects of extreme heat on crop productivity are documented for maize and soybean, which these models were able to capture in a recent study for the U.S.A. (Schauberger *et al* 2017, Anderson *et al* 2015). Another key uncertainty relates to the CO₂ fertilization effect that may lead to enhanced photosynthesis rates and increased crop water productivity, and thereby increased crop productivity under elevated CO₂ concentrations. The strength of this effect is not at all well-constrained by observations and very differently represented in different crop models (Deryng *et al* 2016, see also Fig. 1). (Hasegawa *et al* 2017) suggest that this uncertainty could be reduced for rice, if the reduced effect of CO₂ fertilization on morphological development, in particular leaf area, would be accounted for. This is, however not yet accounted for in the models used here.

In spite of substantial uncertainties in model response, our analysis of crop yield changes at 1.5°C and 2°C for different warming concentration levels indicates that the warming level alone is insufficient to characterise projected impacts of crop productivity. The responsiveness to geophysically plausible CO₂ concentrations at 1.5°C and 2°C is large for most models and crop species and generally outweighs the difference introduced by a half a degree warming increment (Figure 2). This sensitivity remains even for maize, which has no direct CO₂ fertilisation of photosynthesis and only experiences increased water use efficiency under elevated CO₂ (Fig. 1). However, the crop response to elevated CO₂ response in GGCMs has been shown to be a large source of uncertainty (Deryng *et al* 2016) and provides rather optimistic results as models have yet to represent CO₂ interaction processes with, for example, ozone. Another uncertainty dimension relates to the effects of elevated CO₂ on crop quality (Taub *et al* 2008, Myers *et al* 2014), which is a key dimension of food security.

1
2
3 Assessments of climate impacts on crop productivity overlooking the nutrition dimension may easily be
4 misleading with regard to the effect of climate change on future food security (Gustafson *et al* 2016, Müller *et al*
5 2014). Thus, the medium and high CO₂ level scenarios as shown in our study are associated with greater level of
6 uncertainty than our low CO₂ level scenario and should be interpreted with caution.

7 Our analysis highlights consistent negative effects of 0.5°C warming on global and most regional crop
8 productivity for all crops and CO₂ levels investigated. As the climate sensitivity, and thereby the CO₂
9 concentrations at which warming levels of 1.5°C and 2°C may be reached, are inherently uncertain, this has
10 important implications for our understanding of future climate impacts on crop productivity in light of climate
11 sensitivity uncertainty. If TCR turns out to be towards the high end (meaning stronger warming at the same CO₂
12 concentration level), the negative effects of additional warming may subsequently dominate over small (and
13 uncertain) effects of CO₂ fertilization. In the opposite case, stronger CO₂ fertilization, if fully materialized, may
14 dominate, but temperature increase between 1.5°C and 2°C will still lead to adverse impacts (Figure 2). At the
15 same time, a low TCR would allow for a bigger carbon budget to reach warming targets (Rogelj *et al* 2016).
16 Since it is currently not possible to further constrain estimates of TCR, the uncertainty in future impacts on crop
17 productivity under different warming levels is inherently coupled to the geophysical uncertainty of the climate
18 sensitivity (Knutti *et al* 2017).

19 Finally, additional 0.5°C warming increments will consistently lead to more extreme low yields, in particular in
20 tropical regions (Figure 4). Together with a steep rise in world population and food demand over the next
21 decades (Kc and Lutz 2014), this will greatly increase the risk of future food shortages already as early as the
22 2030s when 1.5°C warming could be reached (Lobell and Tebaldi 2014). In a globally connected food system,
23 such production shortages would not only affect the producing regions, but will potentially have strong effects in
24 remote but food importing regions and especially on vulnerable populations that spend large shares of their
25 available income on food. Studies on observed food price shocks linked to extreme weather have indicated that
26 in particular poor, food importing countries – most often least developed countries and small island states – are
27 particularly vulnerable to external production shocks (Bren d'Amour *et al* 2016).

31 32 33 Conclusion

34 Using multi-model multi-ensemble projections for future 1.5°C and 2°C worlds, we have analyzed future
35 changes in crop productivity at these warming levels. We have found consistent negative imprints on 0.5°C
36 warming increments for median as well as low productivity extremes alike for global food productivity with
37 tropical regions being affected more strongly. Despite uncertainties in potential positive effects of elevated CO₂
38 concentrations for crop productivity, we have found that warming levels alone are insufficient to assess future
39 impacts of climate change on future crop productivity. By linking this back to the uncertainty in the geophysical
40 climate response to increased CO₂ emission, our analysis provides a novel viewpoint on the nested geo- and
41 biophysical uncertainties linked to assessments of climate impacts at discrete warming levels. Our findings
42 indicate that impacts of warming on crop production will be consistently lower at 1.5°C compared to 2°C.
43 However, uncertainties related to potentially positive effects of increasing CO₂ fertilization on crop productivity
44 are found to dominate over warming increments. Thereby, our results underscore that GMT levels alone are
45 insufficient to characterise impacts of anthropogenic greenhouse gas emissions on crop productivity.

46 47 48 Acknowledgements

49 The authors would like to thank the HAPPI initiative and all participating modelling groups that have provided
50 data. This research used science gateway resources of the National Energy Research Scientific Computing
51 Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of
52 Energy under Contract No. DE-AC02-05CH11231. CFS and FS acknowledge support by the German Federal
53 Ministry of Education and Research (01LS1613A). DD was supported by the German Federal Ministry for the
54 Environment, Nature Conservation and Nuclear Safety (11_II_093_Global_A_SIDS and LDCs). DD
55 acknowledges the High Performance Computing Cluster supported by the Research and Specialist Computing
56 Support service at the University of East Anglia. WT was supported by an ETH Zurich postdoctoral fellowship
57 (Fel-45 15-1). CM acknowledges financial support from the MACMIT project (01LN1317A) funded through the
58 German Federal Ministry of Education and Research (BMBF). TP acknowledges support from European
59 Commission's 7th Framework Program under Grant Agreement number 603542 (LUC4C). This is paper number
60 36 of the Birmingham Institute of Forest Research. XW acknowledge financial support from AXA.

References

- Anderson C J, Babcock B A, Peng Y, Gassman P W and Campbell T D 2015 Placing bounds on extreme temperature response of maize *Environ. Res. Lett.* **10** 124001
- Asseng S, Ewert F, Martre P, Rötter R, Lobell D, Cammarano D, Kimball B a., Ottman M, Wall G, White J, Reynolds M, Alderman P, Prasad P, Aggarwal P, Anothai J, Basso B, Biernath C, Challinor A, De Sanctis G, Doltra J, Fereres E, Garcia-Vila M, Gayler S, Hoogenboom G, Hunt L, Izaurrealde R, Jabloun M, Jones C, Kersebaum K, Koehler A-K, Müller C, Naresh Kumar S, Nendel C, O'Leary G, Olesen J, Palosuo T, Priesack E, Eyshi Rezaei E, Ruane A, Semenov M, Shcherbak I, Stöckle C, Stratonovitch P, Streck T, Supit I, Tao F, Thorburn P, Waha K, Wang E, Wallach D, Wolf J, Zhao Z and Zhu Y 2014 Rising temperatures reduce global wheat production *Nat. Clim. Chang.* **5** 143–7
- Asseng S, Ewert F and Rosenzweig C 2013 Uncertainty in simulating wheat yields under climate change *Nat. Clim. Chang.* **3** 1–6
- Bren d'Amour C, Wenz L, Kalkuhl M, Christoph Steckel J and Creutzig F 2016 Teleconnected food supply shocks *Environ. Res. Lett.* **11** 35007
- Deryng D, Conway D, Ramankutty N, Price J and Warren R 2014 Global crop yield response to extreme heat stress under multiple climate change futures *Environ. Res. Lett.* **9** 34011
- Deryng D, Elliott J, Folberth C, Müller C, Pugh T A M, Boote K J, Conway D, Ruane A C, Gerten D, Jones J W, Khabarov N, Olin S, Schaphoff S, Schmid E, Yang H and Rosenzweig C 2016 Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity *Nat. Clim. Chang.* **6** 1–7
- Elliott J, Müller C, Deryng D, Chrissyanthacopoulos J, Boote K J, Büchner M, Foster I, Glotter M, Heinke J, Iizumi T, Izaurrealde R C, Mueller N D, Ray D K, Rosenzweig C, Ruane A C and Sheffield J 2015 The Global Gridded Crop Model Intercomparison: Data and modeling protocols for Phase 1 (v1.0) *Geosci. Model Dev.* **8** 261–77
- Frieler K, Betts R, Burke E, Ciais P, Denvil S, Deryng D, Ebi K, Eddy T, Emanuel K, Elliot J, Galbraith E, Gosling S, Halladay K, Hatermann F, Hickler T, Hinkel J, Huber V, Jones C, Krysanova V, Lange S, Lotze H, Lotze-Campen H, Mengel M, Mouratiadou I, Müller Schmied H, Ostberg S, Piontek F, Popp A, Reyer C, Schewe J, Stevanovic M, Suzuki T, Thonicke K, Tian H, Tittensor D T, Vautard R, van Vliet M, Warszawski L and Zhao F 2016 Assessing the impacts of 1.5°C global warming - simulation protocol of the Inter-Sectoral Model Intercomparison Project (ISIMIP2b) *Geosci. Model Dev. Discuss.*
- Greve P and Seneviratne S I 2015 Assessment of future changes in water availability and aridity *Geophys. Res. Lett.* **42** 5493–9
- Gustafson D, Gutman A, Leet W, Drewnowski A, Fanzo J and Ingram J 2016 Seven food system metrics of sustainable nutrition security *Sustain.* **8** 1–17
- Hasegawa T, Li T, Yin X, Zhu Y, Boote K, Baker J, Bregaglio S, Buis S, Confalonieri R, Fugice J, Fumoto T, Gaydon D, Kumar S N, Lafarge T, Marcaida III M, Masutomi Y, Nakagawa H, Oriol P, Ruget F, Singh U, Tang L, Tao F, Wakatsuki H, Wallach D, Wang Y, Wilson L T, Yang L, Yang Y, Yoshida H, Zhang Z and Zhu J 2017 Causes of variation among rice models in yield response to CO₂ examined with Free-Air CO₂ Enrichment and growth chamber experiments *Sci. Rep.* **7** 14858
- IPCC 2014 *Climate Change 2014: Mitigation of Climate Change*. ed O Edenhofer, R Pichs-Madruga, Y Sokona, E Farahani, S Kadner, K Seyboth, A Adler, I Baum, S Brunner, P Eickemeier, B Kriemann, J Savolainen, S Schlömer, C von Stechow, T Zwickel and J C Minx (Cambridge, UK and New York, NY, USA.: Cambridge University Press)
- IPCC 2012 *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation (SREX)* ed C B Field, V Barros, T F Stocker and Q Dahe (Geneva: Cambridge University Press)
- James R, Washington R, Schleussner C-F, Rogelj J and Conway D 2017 Characterizing half-a-degree difference: a review of methods for identifying regional climate responses to global warming targets *Wiley Interdiscip. Rev. Clim. Chang.* e457
- Kc S and Lutz W 2014 The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100 *Glob. Environ. Chang.*
- Kimball B A 2016 Crop responses to elevated CO₂ and interactions with H₂O, N, and temperature *Curr. Opin. Plant Biol.* **31** 36–43
- Knutti R, Rugenstein M A A and Hegerl G C 2017 Beyond equilibrium climate sensitivity *Nat. Geosci.*
- Lange S 2017 Bias correction of surface downwelling longwave and shortwave radiation for the EWEMBI dataset *Earth Syst. Dyn. Discuss.* **2017** 1–30
- Lesk C, Rowhani P and Ramankutty N 2016 Influence of extreme weather disasters on global crop production *Nature* **529** 84–7
- Liu B, Asseng S, Müller C, Ewert F, Elliott J, Lobell D B B, Martre P, Ruane A C C, Wallach D, Jones J W W,

- Rosenzweig C, Aggarwal P K K, Alderman P D D, Anothai J, Basso B, Biernath C, Cammarano D, Challinor A, Deryng D, Sanctis G D De, Doltra J, Fereres E, Folberth C, Garcia-Vila M, Gayler S, Hoogenboom G, Hunt L A A, Izaurralde R C C, Jabloun M, Jones C D D, Kersebaum K C C, Kimball B A A, Koehler A-K, Kumar S N N, Nendel C, O'Leary G G J, Olesen J E E, Ottman M J J, Palosuo T, Prasad P V V V V, Priesack E, Pugh T A M A M, Reynolds M, Rezaei E E E, Rötter R P P, Schmid E, Semenov M A A, Shcherbak I, Stehfest E, Stöckle C O O, Stratonovitch P, Streck T, Supit I, Tao F, Thorburn P, Waha K, Wall G W W, Wang E, White J W J W W, Wolf J, Zhao Z and Zhu Y 2016 Similar estimates of temperature impacts on global wheat yield by three independent methods *Nat. Clim. Chang.* **1** 1–8
- Lobell D B and Asseng S 2017 Comparing estimates of climate change impacts from process- based and statistical crop models *Environ. Res. Lett.* **12** 1–12
- Lobell D B, Bänziger M, Magorokosho C and Vivek B 2011a Nonlinear heat effects on African maize as evidenced by historical yield trials *Nat. Clim. Chang.* **1** 42–5
- Lobell D B, Schlenker W and Costa-Roberts J 2011b Climate trends and global crop production since 1980. *Science* **333** 616–20
- Lobell D B and Tebaldi C 2014 Getting caught with our plants down: the risks of a global crop yield slowdown from climate trends in the next two decades *Environ. Res. Lett.* **9** 74003
- McGrath J M and Lobell D B 2013 Regional disparities in the CO₂ fertilization effect and implications for crop yields *Environ. Res. Lett.* **8** 14054
- Mitchell D, AchutaRao K, Allen M, Bethke I, Beyerle U, Ciavarella A, Forster P M, Fuglestedt J, Gillett N, Haustein K, Ingram W, Iversen T, Kharin V, Klingaman N, Massey N, Fischer E, Schleussner C-F, Scinocca J, Seland Ø, Shiogama H, Shuckburgh E, Sparrow S, Stone D, Uhe P, Wallom D, Wehner M and Zaaboul R 2017 Half a degree additional warming, prognosis and projected impacts (HAPPI): background and experimental design *Geosci. Model Dev.* **10** 571–83
- Moore F C and Lobell D B 2015 The fingerprint of climate trends on European crop yields *Proc. Natl. Acad. Sci.* 201409606
- Morgan J A, LeCain D R, Pendall E, Blumenthal D M, Kimball B A, Carrillo Y, Williams D G, Heisler-White J, Dijkstra F A and West M 2011 C4 grasses prosper as carbon dioxide eliminates desiccation in warmed semi-arid grassland *Nature* **476** 202–5
- Müller C, Elliott J, Chryssanthacopoulos J, Arneth A, Balkovic J, Ciais P, Deryng D, Folberth C, Glotter M, Hoek S, Iizumi T, Izaurralde R C, Jones C, Khabarov N, Lawrence P, Liu W, Olin S, Pugh T A M, Ray D K, Reddy A, Rosenzweig C, Ruane A C, Sakurai G, Schmid E, Skalsky R, Song C X, Wang X, de Wit A and Yang H 2017 Global gridded crop model evaluation: benchmarking, skills, deficiencies and implications *Geosci. Model Dev.* **10** 1403–22
- Müller C, Elliott J, Chryssanthacopoulos J, Deryng D, Folberth C, Pugh T A M and Schmid E 2015 Implications of climate mitigation for future agricultural production *Environ. Res. Lett.* **10** 125004
- Müller C, Elliott J and Levermann A 2014 Food security: Fertilizing hidden hunger *Nat. Clim. Chang.* **4** 540–1
- Myers S S, Zanobetti A, Kloog I, Huybers P, Leakey A D B, Bloom A J, Carlisle E, Dieterich L H, Fitzgerald G, Hasegawa T, Holbrook N M, Nelson R L, Ottman M J, Raboy V, Sakai H, Sartor K A, Schwartz J, Seneweera S, Tausz M and Usui Y 2014 Increasing CO₂ threatens human nutrition. *Nature* **510** 139–42
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000-Global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling *Global Biogeochem. Cycles* **24**n/a-n/a
- Porwollik V, Müller C, Elliott J, Chryssanthacopoulos J, Iizumi T, Ray D K, Ruane A C, Arneth A, Balković J, Ciais P, Deryng D, Folberth C, Izaurralde R C, Jones C D, Khabarov N, Lawrence P J, Liu W, Pugh T A M, Reddy A, Sakurai G, Schmid E, Wang X, de Wit A and Wu X 2017 Spatial and temporal uncertainty of crop yield aggregations *Eur. J. Agron.* **88** 10–21
- Ray D K, Gerber J S, Macdonald G K and West P C 2015 Climate variation explains a third of global crop yield variability *Nat. Commun.* **6** 1–9
- Rogelj J, Schaeffer M, Friedlingstein P, Gillett N P, van Vuuren D P, Riahi K, Allen M and Knutti R 2016 Differences between carbon budget estimates unravelled *Nat. Clim. Chang.* **6** 245–52
- Rosenzweig C, Elliott J, Deryng D, Ruane A C, Müller C, Arneth A, Boote K J, Folberth C, Glotter M, Khabarov N, Neumann K, Piontek F, Pugh T a M, Schmid E, Stehfest E, Yang H and Jones J W 2014 Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison *Proc. Natl. Acad. Sci.* **111** 3268–73
- Schauberger B, Archontoulis S, Arneth A, Balkovic J, Ciais P, Deryng D, Elliott J, Folberth C, Khabarov N, Müller C, Pugh T A M, Rolinski S, Schaphoff S, Schmid E, Wang X, Schlenker W and Frieler K 2017 Consistent negative response of US crops to high temperatures in observations and crop models *Nat. Commun.* **8** 13931
- Schlenker W and Lobell D B 2010 Robust negative impacts of climate change on African agriculture *Environ. Res. Lett.* **5** 14010
- Schleussner C-F, Lissner T, Fischer E, Wohland J, Perrette M, Golly A, Rogelj J, Childers K, Schewe J, Frieler K, Hare W and Schaeffer M 2016a Differential climate impacts for policy relevant limits to global

- 1
2
3 warming: the case of 1.5°C and 2°C *Earth Syst. Dyn.* **7** 327–51
- 4 Schlessner C-F, Rogelj J, Schaeffer M, Lissner T, Licker R, Fischer E M, Knutti R, Levermann A, Frieler K
5 and Hare W 2016b Science and policy characteristics of the Paris Agreement temperature goal *Nat. Clim.*
6 *Chang.* **6** 827–835
- 7 Stocker T F, Qin D, Plattner G-K, Alexander L V, Allen S K, Bindoff N L, Bréon F-M, Church J A, Cubasch U,
8 Emori S, Forster P, Friedlingstein P, Gillett N, Gregory J M, Hartmann D L, Jansen E, Kirtman B, Knutti
9 R, Kumar K K, Lemke P, Marotzke J, Masson-Delmotte V, Meehl G A, Mokhov I I, Piao S, Ramaswamy
10 V, Randall D, Rhein M, Rojas M, Sabine C, Shindell D, Talley L D, Vaughan D G and Xie S-P 2013
11 Technical Summary *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
12 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker, D Qin,
13 G-K Plattner, M Tignor, S K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (Cambridge,
14 UK and New York, NY, USA.: IPCC AR5 WGI, Cambridge University Press)
- 15 Taub D R, Miller B and Allen H 2008 Effects of elevated CO₂ on the protein concentration of food crops: A
16 meta-analysis *Glob. Chang. Biol.* **14** 565–75
- 17 Thiery W, Davin E L, Lawrence D M, Hirsch A L, Hauser M and Seneviratne S I 2017 Present-day irrigation
18 mitigates heat extremes *J. Geophys. Res. Atmos.* **122** 1403–22
- 19 UNFCCC 1992 *United Nations Framework Convention on Climate Change* (UNFCCC)
- 20 Wang E, Martre P, Zhao Z, Ewert F, Maiorano A, Rötter R P, Kimball B A, Ottman M J, Wall G W, White J W,
21 Reynolds M P, Alderman P D, Aggarwal P K, Anothai J, Basso B, Biernath C, Cammarano D, Challinor A
22 J, De Sanctis G, Doltra J, Fereres E, Garcia-Vila M, Gayler S, Hoogenboom G, Hunt L A, Izaurralde R C,
23 Jabloun M, Jones C D, Kersebaum K C, Koehler A K, Liu L, Müller C, Naresh Kumar S, Nendel C,
24 O’Leary G, Olesen J E, Palosuo T, Priesack E, Eyshi Rezaei E, Ripoche D, Ruane A C, Semenov M A,
25 Shcherbak I, Stöckle C, Stratonovitch P, Streck T, Supit I, Tao F, Thorburn P, Waha K, Wallach D, Wang
26 Z, Wolf J, Zhu Y and Asseng S 2017 The uncertainty of crop yield projections is reduced by improved
27 temperature response functions *Nat. Plants* **3**
- 28 Wang J, Wang C, Chen N, Xiong Z, Wolfe D and Zou J 2015 Response of rice production to elevated [CO₂] and
29 its interaction with rising temperature or nitrogen supply: a meta-analysis *Clim. Change* **130** 529–43
- 30 Warszawski L, Frieler K, Huber V, Piontek F, Serdeczny O and Schewe J 2013 The Inter-Sectoral Impact Model
31 Intercomparison Project (ISI-MIP): Project framework. *Proc. Natl. Acad. Sci. USA* 1–5
- 32 Zhao C, Liu B, Piao S, Wang X, Lobell D B, Huang Y, Huang M, Yao Y, Bassu S, Ciais P, Durand J-L, Elliott
33 J, Ewert F, Janssens I A, Li T, Lin E, Liu Q, Martre P, Müller C, Peng S, Peñuelas J, Ruane A C, Wallach
34 D, Wang T, Wu D, Liu Z, Zhu Y, Zhu Z and Asseng S 2017 Temperature increase reduces global yields of
35 major crops in four independent estimates *Proc. Natl. Acad. Sci.* **114** 9326–31
- 36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript