



Mitigation pathways of air pollution from residential emissions in the Beijing-Tianjin-Hebei region in China



Jun Liu^{a,b,*}, Gregor Kiesewetter^a, Zbigniew Klimont^a, Janusz Cofala^a, Chris Heyes^a, Wolfgang Schöpp^a, Tong Zhu^c, Guiying Cao^a, Adriana Gomez Sanabria^a, Robert Sander^a, Fei Guo^a, Qiang Zhang^b, Binh Nguyen^a, Imrich Bertok^a, Peter Rafaj^a, Markus Amann^{a,**}

^a International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, A-2361 Laxenburg, Austria

^b Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing 100084, China

^c State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

ARTICLE INFO

Handling Editor: Xavier Querol

Keywords:

Solid fuel substitution

PM_{2.5}

Natural gas

Human exposure

Outdoor and indoor health benefits

ABSTRACT

Air pollution is one of the most harmful consequences of China's rapid economic development and urbanization. Particularly in the Beijing-Tianjin-Hebei (BTH) regions, particulate matter concentrations have consistently exceeded the national air quality standards. Over the last years, China implemented ambitious measures to reduce emissions from the power, industry and transportation sectors, with notable success during the 11th and 12th Five Year Plan (FYP) periods. However, such strategies appear to be insufficient to reduce the ambient PM_{2.5} concentration below the National Air Quality Standard of 35 $\mu\text{g m}^{-3}$ across the BTH region within the next 15 years. We find that a comprehensive mitigation strategy for the residential sector in the BTH region would deliver substantial air quality benefits. Beyond the already planned expansion of district heating and natural gas distribution in urban centers and the foreseen curtailment of coal use for households, such a strategy would redirect some natural gas from power generation units towards the residential sector. Rural households would replace biomass for cooking by liquid petroleum gas (LPG) and electricity, and substitute coal for heating by briquettes. Jointly, these measures could reduce the primary PM_{2.5} and SO₂ emissions by 28% and 11%, respectively, and the population-weighted PM_{2.5} concentrations by 13%, i.e., from 68 $\mu\text{g m}^{-3}$ to 59 $\mu\text{g m}^{-3}$. We estimate that such a strategy would reduce premature deaths attributable to ambient and indoor air pollution by almost one third.

1. Introduction

Air pollution is one of the most harmful consequences of China's rapid economic development and urbanization. Over large areas, ambient concentrations of PM_{2.5} (particulate matter with aerodynamic diameter equal to or smaller than 2.5 μm) exceed the Air Quality Guidelines of the World Health Organization (WHO) as well as National Air Quality Standards. Exposure of the population to such high pollution levels causes a significant burden to public health and economic development, although precise quantification remains uncertain. Published estimates range between 0.35 million (Chen et al., 2013) and more than 1.1 million cases (Cohen et al., 2017) of premature death annually that are attributable to outdoor air pollution in China, and indicate even larger impacts on morbidity. Associated welfare losses

have been valued at between 3.8% of GDP (World Bank, 2007) and 9.9% of GDP (World Bank, 2016).

Due to heavy industrialization, high population density and meteorological conditions, the Beijing-Tianjin-Hebei (BTH in short) region in the Northern China Plain is one of the most polluted hotspots in China, next to the Yangtze River Delta and the Pearl River Delta. During 2013-2014, monitoring stations located in cities in the BTH region measured the highest annual average ambient PM_{2.5} concentration in China - up to 144 $\mu\text{g m}^{-3}$ in Shijiazhuang, followed by Tianjin with 92 $\mu\text{g m}^{-3}$, and Beijing with 87 $\mu\text{g m}^{-3}$ (Wang et al., 2014a). Thereby, annual average ambient PM_{2.5} concentrations in the BTH region were up to four times higher than China's Ambient Air Quality Standard level II of 35 $\mu\text{g m}^{-3}$ (equivalent to the Interim Target Level 1 of the World Health Organization [WHO]).

* Correspondence to: J. Liu, International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, A-2361 Laxenburg, Austria.

** Corresponding author.

E-mail addresses: liujun@iiasa.ac.at (J. Liu), amann@iiasa.ac.at (M. Amann).

In response, the Chinese government has established ambitious policy targets to improve air quality and initiated comprehensive plans and programs to achieve these targets. In contrast to most other countries in the world, in China the largest share of pollution used to originate from power generation and industry, caused, inter alia, by the heavy reliance on coal (Wang et al., 2012; Zhang et al., 2012). Consequently, the emission reduction programs used to focus on these sectors. However, the contributions of other sources, including households and agricultural activities, are sizeable as well, even if they did not receive the same policy attention in the past (Liu et al., 2016; Zhao et al., 2017). Only recently, the authorities started to address mitigation in the residential sector, and are striving to switch fuel use in households (MEP, 2017).

In this study, we employ the Greenhouse gas – Air pollution Interactions and Synergies (GAINS) model (Amann et al., 2011) to facilitate a systematic integrated exploration of effective policy intervention options. By tracking the pathways of air pollutants from energy consumption to emissions, the atmospheric chemistry and transport processes, and the human health impacts, we develop scenarios to seek opportunities in residential clean heating and cooking in the BTH region during 2010–2030. The study explores mitigation pathways for the residential sector, and shows that re-thinking of the energy supply strategy could deliver large benefits on air quality and development, and make an efficient contribution to China's endeavor to achieve cleaner air.

2. Methodology

2.1. GAINS model

The scenarios are built on IIASA's (International Institute for Applied Systems Analysis) Greenhouse gas – Air pollution Interactions and Synergies (GAINS) model. As a scientific tool for integrated policy assessment, the GAINS model describes the air pollution pathways from atmospheric driving force to environmental impact, with information on economic, energy and agricultural development, emission control strategies and cost, atmospheric dispersion and source sensitivities (Amann et al., 2011). It quantifies the emissions and impacts of six air pollutants (SO₂, NO_x, CO, NH₃, VOCs, PM_{2.5}, PM_{2.5-10}, PM₁, BC, OC) and six greenhouse gases (CO₂, CH₄, N₂O, HFCs, PFCs, SF₆) on human health, crop losses, acid deposition, and long-term radiative forcing in a multi-pollutant perspective. This interdisciplinary model has already guided European governments in key negotiations on air pollution control agreements across the continent without compromising economic development (Amann et al., 2008; Amann et al., 2015). It has been applied for analysis of air pollutant emissions and air quality and health impacts in the IEA 2016 World Energy Outlook Special Report on Energy and Air Pollution (IEA, 2016), and has also been developed to evaluate co-benefits of emission reductions on the city scale (GAINS-City) (Amann et al., 2017; Liu et al., 2013).

The GAINS model estimates emissions with technology-based methodology following equation (Amann et al., 2011):

$$E_p = \sum_k \sum_m A_k e_{f_{k,m,p}} \chi_{k,m,p} \quad (1)$$

where, k , m , p represents activity type, abatement measure, pollutant respectively; E_p stands for emission of pollutant p , A_k stands for activity data of type k , $e_{f_{k,m,p}}$ stands for emission factor of pollutant p for activity k after application of control measure m , $\chi_{k,m,p}$ stands for penetration of control measure m for pollutant p of activity k . The emission factors used in the GAINS model for Asia were originally collected within the GAINS-Asia project (Amann et al., 2008). The current version of the model incorporates updates, which use literature reported measurements of emissions from residential combustion sources in China (Cao et al., 2008; Chen et al., 2009; Li et al., 2009; Zhi et al., 2008, 2009) as well as local information on parameters to calculate

emission factors from large stationary combustion sources including coal-fired power plants. For instance, updates were made for sulphur and ash content of fuels, as well as efficiency and penetration rates of control measures (Klimont et al., 2009, 2017; Zhao et al., 2013). These emission factors have been applied to estimate air pollutant emissions for the World Energy Outlook Special Report 2016: Energy and Air pollution (IEA, 2016). The PM_{2.5}, SO₂ and NO_x emission factors used in this study are summarized in Table S1.

2.2. Emission scenarios

Two scenarios are constructed and analyzed in this study: Baseline and Household scenario. In the Baseline, fuel consumption by sector and by province in base year 2010 is obtained from the China Energy Statistical Yearbook (National Bureau of Statistics, 2011), and is implemented in the GAINS model as the base year activity. The GAINS approach and, in particular, the methodology for the household and commercial sector is described in Klimont et al. (2017). Quantities of fuels were converted into energy equivalents using net calorific values (Table S2) in the GAINS model. The household energy consumption for cooking is obtained from recent Chinese household surveys (Table S3) (Liang et al., 2013; Zheng et al., 2014), and the residues from the total household energy consumption are distributed as fuel for heating. Table S4 presents the resulting residential energy consumption by fuel type and final application in the BTH region in the base year 2010.

For the future scenarios, an annual economic growth of 6% per year up to 2030 is assumed, following the projections of the IEA World Energy Outlook 2015 (IEA, 2015), which is in accordance with China's recent economic growth and the economic development target in the Thirteenth Five-year Plan. A continuing urbanization (2%/a) was assumed based on the government development plan of the three regions (Beijing Municipal Government, 2017; Tianjin Municipal Government, 2006; DHUDH, 2017), and the population growth trend was forecast by the IIASA world population projection model (Lutz et al., 2014), based on a moderate change of fertility, mortality, and inter-provincial migration up to 2030 in Beijing, Tianjin and Hebei. Energy use projections are based on the provincial energy use pattern in 2010 from the China Energy Statistical Yearbook and the national trends from the International Energy Agency (IEA) World Energy Outlook 2015 (IEA, 2015), with additional assumptions on the evolution of energy consumption for individual use in the residential sector to reflect current policies of limiting residential coal and biomass use in the BTH region. The development of other fuel consumption (LPG, heat, renewable energy, liquid fuel and electricity) is consistent with the trend from the IEA WEO (2015) projection, and the natural gas demand is determined by meeting the assumptions for energy consumption for individual needs and coal consumption decrease. Details on the policy incorporation and the establishment of the Baseline scenario are provided in the Supporting information (SI).

Starting from the Baseline, the Household scenario targeting emission reductions from cleaner fuel use in the household and commercial sectors employs measures on four aspects: 1) in the commercial sector, all the solid use is replaced by natural gas; 2) in the urban and suburban areas, all the solid fuel for heating is completely replaced by natural gas; similarly, all LPG for cooking is substituted with natural gas; 3) the substituted LPG for cooking in the urban and suburban areas is used to replace coal and biomass for cooking in rural areas, and to completely phase out solid fuels for cooking in rural areas, additional wind power is used to replace the remaining coal and biomass for cooking; 4) considering the potentially high distribution cost of natural gas due to low population density, the demand for solid fuel for heating in rural areas will not be replaced with natural gas, but will be supplied with coal briquettes and biomass briquettes instead. Details on the building of the Household scenarios and methods for fuel substitution are provided in the SI.

2.3. Modelling PM_{2.5} concentration and health impact estimation

The air quality in each scenario is estimated with the GAINS model. The annual ambient PM_{2.5} concentration in GAINS includes two components (Kiesewetter et al., 2015): 1) a regional background concentration related to emissions (including PM_{2.5}, SO₂, NO_x, NH₃ and VOCs) from Chinese provinces and other Asian countries calculated using linear source-receptor transfer coefficients derived from sensitivity simulations of the EMEP (European Monitoring and Evaluation Programme) chemical transport model (Simpson et al., 2012) at a resolution of 0.5° × 0.5°; 2) a sub-grid increment related to primary PM_{2.5} emissions from local, low-level sources (including residential and transportation emissions), with grid-specific local, low-level PM_{2.5} transfer coefficients derived from linear regression analysis of the low-level emission EMEP model perturbation runs. For the calculation of the sub-grid increment, to make the calculations manageable, the cities with over 100,000 inhabitants identified by Global Rural-Urban Mapping Project (GRUMP) v1 dataset are defined as large urban areas in the model, while the remaining smaller cities and other remaining regions are defined as suburban and rural areas, in order to distribute the urban and rural low-level primary PM_{2.5} emissions separately. Therefore, the PM_{2.5} concentrations in small cities may be underestimated if the low-level primary PM_{2.5} emission intensity in small cities is higher than that in rural areas within the same grid cell. Then the combined results are used to estimate population exposure of ambient PM_{2.5} concentrations. Details on the methodology for the air quality modelling are provided in the SI.

Since 2013 the Chinese national monitoring network has included direct measurements of PM_{2.5}, and the observed PM_{2.5} annual average concentrations for major cities are obtained from the China Statistical Yearbook (National Bureau of Statistics, 2016). For the base year 2010, there are limited PM_{2.5} observations available in China, so model validation is carried out with the observed PM_{2.5} annual concentrations for 2015 (Fig. S1 and Table S5). The modelled annual PM_{2.5} concentrations agree well with observations, with correlation coefficients at 0.8, and RMSE at 16.3 μg m⁻³. 98% of the modelled concentrations are within a factor of two of the observations, although the model slightly overestimates observations (NMB 5%). Overall, the model evaluation results are comparable with other model studies over China (Geng et al., 2017; Ma et al., 2017).

For health impacts, we estimate the premature mortality attributable to PM_{2.5} exposure from both ambient air pollution and indoor air pollution for each scenario. The exposure to ambient air pollution is calculated by the overlay of modelled annual average PM_{2.5} concentrations and population distribution. The household air pollution is calculated for residents using solid fuel cooking. A typical exposure of 300 μg m⁻³ is assumed for traditional stoves and 70 μg m⁻³ for clean stoves, based on the wide range of observed concentrations (Balakrishnan et al., 2013). The human health impact in terms of premature mortalities of each scenario is estimated for all relevant diseases, including ischemic heart disease (IHD), cerebrovascular disease (stroke), chronic obstructive pulmonary disease (COPD), and lung cancer (LC) in five-year age classes. Baseline mortalities are taken from the GBD-2013 database. The integrated exposure–response (IER) function (Burnett et al., 2014) developed for the GBD-2013 study (Forouzanfar et al., 2015) and used in the WHO 2016 assessment of ambient air pollution (WHO, 2016) is used to estimate the relative risk associated with a certain level of PM_{2.5} exposure. Details on the methodology for the air quality modelling and health impact estimation are provided in the SI.

3. Results

3.1. Air pollution in the BTH region

For this study, we employ the East-Asia implementation of the

GAINS model, which distinguishes 31 regions and provinces in China, including the BTH region. Relying on provincial energy statistics, fuel characteristics and emission inventories, we estimate with the GAINS model that in 2010 power plants and industry were the largest sources of SO₂, NO_x and primary PM_{2.5} emissions in BTH, responsible for 85%, 66% and 52% of total emissions, respectively (Table S6). Transportation caused 30% of NO_x and 5% of PM_{2.5} emissions. In comparison, the residential sector (defined as households and commercial sector in this study) was a smaller source, with 35% of total PM_{2.5}, 15% of SO₂ and 5% of NO_x emissions. Agricultural activities (manure management and mineral nitrogen fertilizer application, primarily urea) made the predominant contributions (more than 90%) to NH₃ emissions in the region.

However, stack heights and proximities between emissions and population differ across these sectors. Based on emission inventories and the EMEP atmospheric chemistry and transport model, GAINS quantifies the contributions of the various source categories to ambient concentrations of PM_{2.5} across the BTH region with a resolution of 0.5° × 0.5° (roughly 40 × 50 km), on top of which concentrations are calculated explicitly at higher resolution for each individual city with more than 100,000 inhabitants. Combined with population density data, the total population exposure to PM_{2.5} across the BTH region can then be computed.

Using such a methodology, we estimate for the BTH area a population-weighted PM_{2.5} concentration of 85 μg m⁻³ in 2010. For 96% of the population, exposure of PM_{2.5} exceeded the China Ambient Air Quality level II of 35 μg m⁻³. 79% of the people faced PM_{2.5} levels of more than twice this level, 15% more than three times, and about 3% more than four times (Fig. 1).

Source attributions suggest that, averaged for the entire population within the BTH region, about 60% of the exposure to PM_{2.5} in outdoor air originates from emissions in the BTH region, and another 34% from other Chinese provinces outside BTH. The contribution from natural sources (soil dust, sea salt) is estimated at 6% (Fig. 2A). Similar findings are proposed in previous studies (Li et al., 2015; An et al., 2007). Considering the BTH contributions, about one-third is the local concentration increment contributed by low-level residential and transportation primary PM_{2.5} emissions (Fig. 2A), estimated with the downscaling methodology. The share of local primary concentration is even higher for Hebei conurbations, where the residential emissions are higher (Fig. 2B). From the population exposure that is caused by emission sources within the BTH area (60% of total exposure), the largest contribution comes from the power and industrial sectors (25%), 18% from residential sources (i.e., burning of coal and biomass for heating and cooking), 11% from agricultural emissions, 4% from transportation, and 2% from waste burning sources in the BTH region (Fig. 2A).

3.2. Outlook to 2030

The rapid economic development and urbanization, together with the re-orientation of industrial and energy policies and the ambitious sharpening of pollution control legislation and enforcement, will impact future pollution levels in China. We use the GAINS model to assess the likely interplay of these factors on future air quality in the BTH region, identify the most important pollution sources that will remain after implementation of current policies, and explore policy intervention options from small combustion sources in the residential and commercial sector.

3.2.1. Economic development, energy policy and pollution controls

Following the central projections of fertility, mortality and inter-provincial migration of the integrated development plan of Beijing, Tianjin and Hebei in the next 15 years, we project population in BTH to increase by 21% compared to 2010. Urbanization will reach 78%, with a 68% increase of urban and a 40% decline of rural population (Table




Fig. 1. Ambient PM_{2.5} concentrations (A) and proportion of population exposed to different levels of PM_{2.5} (B) in the BTH region in 2010, computed with GAINS model.




Fig. 2. Source apportionment for population-weighted PM_{2.5} concentration in the BTH region (A) and Hebei conurbations (B) in 2010. The source regions are distinguished by regions outside of BTH and regions within BTH; five types of sources are identified, including natural, power plus industry, residential, transportation, agriculture, waste burning, and international shipping. The contribution of primary PM_{2.5} increment from local low-level residential and transportation sources are estimated individually with the down-scaling method.

S7). For economic development and energy consumption we adopt the scenario of the IEA as presented in its World Energy Outlook 2016 Special Report on ‘Energy and Air Pollution’ (IEA, 2016), which assumes a continued economic growth of 6%/year. This will triple per capita GDP in 2030.

The projection assumes continuation of the ongoing trend towards consumption and services, and further transformation towards less energy-intensive production. Together with the latest policy measures, this will reduce the energy intensity of GDP by 54%, although total primary energy use will still increase by 54%. Overall, the share of coal in the primary energy mix of the BTH region will decline from about two-thirds in 2010 to 50% in 2030, although the absolute volume will still increase by 25%. Natural gas will double its share, and consumption will grow by almost a factor of two. 26% of the natural gas volume will be consumed in the commercial sector and households, while the amount of natural gas distributed in the power and industrial sectors is larger, accounting for 34% and 37% of the total supply, respectively.

Growing population and economic wealth will increase energy consumption in the households and commercial sector by 50%. At the same time, we assume that the latest policies on limiting residential coal and biomass use in the BTH region will decrease solid fuel consumption for heating in urban areas by 26%, and by 44% in the rural area following the decline in rural population. For cooking, we assume that by 2030 all solid fuel use will disappear in all urban centers with

population densities of more than 20 people/ha, and remain only in the urban fringes, suburban areas, small towns and rural areas (Table S8).

Although past experience with respect to compliance with legislation has not always been positive, we assume full implementation and effective enforcement of the measures and policies that have been decided by 2015. These include the measures outlined in the 12th Five-year Plan of the Chinese Government and more specifically in the Air Pollution Prevention and Control in Key Areas (MEP, 2012), as well as the measures specified in the 2013 Air Pollution Prevention and Control Action Plan of the State Council (hereafter referred to as the ‘Action Plan’) (MEP, 2013), and the Implementation Plan specially for BTH and its surrounding regions. However, we consider only the specific pollution control measures outlined in these documents, but do not speculate about additional measures that might be required to fully achieve China’s targets on ambient air quality. Furthermore, as at the time of writing there is no available translation of the 13th Five Year Plan into specific measures, we do not consider additional measures that might emerge from this document.

3.2.2. Emissions and population exposure

The policy measures outlined above will achieve a distinct decoupling between economic growth, energy use and pollution levels in the BTH region. Despite the doubling of GDP and a 50% higher energy consumption in 2030, SO₂, NO_x and PM_{2.5} emissions would drop by

Table 1
BTH air pollutant and greenhouse gas emissions, population-weighted PM_{2.5} concentration, and human health impacts in 2010 and 2030 for each scenario.

	2010	2030 Baseline scenario	2030 Household scenario
PM _{2.5} (kt)	1160	728	524
SO ₂ (kt)	2483	1393	1235
NO _x (kt)	2441	1673	1634
NH ₃ (kt)	1059	1196	1192
CO ₂ (Mt)	944	1265	1261
CH ₄ (Mt CO ₂ eq)	60.8	78.0	77.8
PM _{2.5} population-weighted exposure (μg m ⁻³)	85	68	59
Households (population) using solid fuels for cooking	14.0 (42.6) million	7.3 (24.2) million	0
Premature deaths			
From ambient pollution	89,788	118,742	113,250
From household air pollution caused by solid fuel use for cooking	73,865	50,880	0

44%, 31% and 37%, respectively, as a result of the new policies (Table 1). However, in the absence of any emission regulation for the agricultural sector, NH₃ emissions, an important precursor of PM_{2.5}, will grow by 13% following closely the production trend.

Assuming that similar policies will be implemented in the provinces surrounding BTH, population-weighted PM_{2.5} concentrations in the BTH region would decline from 85 μg m⁻³ in 2010 to 68 μg m⁻³ in 2030, i.e., by 20% (Table 1). As a consequence, fewer people will be exposed to extreme levels; e.g., while in 2010 almost 80% of the population faced more than 70 μg m⁻³ of PM_{2.5} (i.e., more than double the Chinese Ambient Air Quality standard II and the WHO Interim Target Level 1), this share will drop to 45% in 2030 (Fig. S2). In addition, 24.2 million people who were using solid fuels for cooking and heating will switch to clean fuels, and will be thereby relieved from the health burden of indoor pollution caused by solid fuel use for cooking.

However, these reductions will not be sufficient to significantly enlarge the share of population facing concentrations that comply with the current Chinese air quality standards. We estimate major contributions from natural sources and emitters outside the BTH region (about 25 μg m⁻³ on a population-weighted basis). Residual emissions from the industrial and power sectors in BTH will still contribute about 16 μg m⁻³, the household and commercial sector 11 μg m⁻³, agricultural activities 11 μg m⁻³, and other sources including transportation about 5 μg m⁻³ (Fig. S2). It should be noted that further efforts from the Acton Plan and 13th FYP are underway, which will significantly reduce industrial emissions, especially SO₂, and the 16 μg m⁻³ contribution of industrial and power sectors can be further reduced by these latest efforts (Zheng et al., 2018).

3.3. An alternative strategy for household energy use

3.3.1. The household scenario

While, for good reasons, significant attention has been paid to pollution controls in power plants and industry in China, the residential sector, despite making the second-largest contribution to population exposure in 2010, has received less consideration. Per unit of energy use, solid fuel combustion in small, low-level sources in the household and commercial sector has a significantly stronger impact on ambient air quality and population exposure than large point sources, and its strong seasonality makes it a key source in pollution episodes during the heating season, as confirmed by source apportionment (Yang et al., 2016; Li et al., 2015) and remote sensing (Xiao et al., 2015) studies. According to the estimates presented in Section 3.2.2, in 2030 solid fuel use in the residential sector of the BTH region will still account for 16% of the population exposure (including sources outside the BTH region),

despite the measures taken in the baseline scenario.

It is noteworthy that in spite of the policies for household energy use, (i.e., expansion of district heating, phase-out of coal for heating and cooking in urban centers, etc.), in 2030 a substantial amount of solid biomass and coal remains for cooking and heating in suburban and rural areas. At the same time, natural gas use will be expanded by 170%; however, only less than one-third of the additional gas volume is directed to the residential sector, while more than two-thirds will be used to substitute coal in large industrial installations and power plants.

As there are only limited possibilities to reduce emissions from solid fuel use in the residential sector without fuel switches, while there are ample opportunities for controlling emissions at large installations through end-of-pipe technologies, we explore in an alternative scenario the practical scope for air quality improvements in the BTH region from an enhanced penetration of clean fuels to the residential sector.

In 2015, total heated floor space in the cities of Hebei amounted to 1.07 billion m², 80% connected to central heating systems (HPDHURC, 2017). However, despite major investments into grid-based energy supply in the cities, there remains a persistent fraction of coal use, mainly among low-skilled rural-to-urban migrants due to housing conditions, easy access, and low costs of coal (Ru et al., 2015). In the entire BTH region, estimated with GAINS, district heating supplied 56% of the demand in cities and sub-urban areas, other forms of clean energy (natural gas, electricity and renewable energy) about 13%, and about 29% of the population is still using coal for heating.

For the future, we assume for the urban areas in the BTH region that all coal and biomass for heating, as well as coal for the commercial sector, will be completely replaced by natural gas. Similarly, all LPG used for cooking in urban areas will be shifted to rural areas to substitute solid fuels for cooking. Furthermore, rural households that are currently using solid fuels for cooking will switch to electric stoves, with the electricity generated by additional wind power. This will increase wind power generation by 13% above the baseline projection, a small margin compared to the seven-fold growth that is assumed there (China Electric Power Yearbook Editorial Committee, 2011, 2015; IEA, 2015). Furthermore, as a conservative assumption, we do not consider replacement of solid fuels for heating in rural areas by natural gas due to potentially high distribution costs in low-density areas. Instead, the residual demand for solid fuels for heating in rural areas will be supplied by coal briquettes and biomass briquettes.

Such a strategy would increase the projected demand for natural gas in the residential sector by 21% (or 5% of total gas demand in the BTH region) compared to the baseline (Table S9). Most likely, this additional quantity could be obtained on the market (e.g., additional imports from Russia, shale gas, LNG). However, as a conservative assumption, we divert this gas volume from the power sector and replace it there by coal, which will increase coal consumption in the power sector by 6%.

Compared to the baseline scenario, such a strategy would replace in urban areas (mainly suburban) 11.3 million tons of coal and 0.9 million tons of biomass used for heating by 4.6 bcm of natural gas. In 4.2 million (sub-)urban households, 0.69 million tons of LPG used for cooking will be substituted by 0.84 bcm of natural gas. This will allow 3.4 million households in rural areas to replace biomass and coal for cooking by LPG, while 3.9 million rural households will switch to electricity for cooking. For heating, 6.1 million rural households will replace conventional coal and biomass with briquettes. In total, this would reduce biomass combustion in the BTH area by 29% and coal consumption by 2% compared to the baseline; natural gas use will remain the same, and wind power increase by 13%.

3.3.2. Emissions, air quality and health impacts

Such a re-arrangement of fuel use will have major beneficial impacts on air pollutant emissions: in 2030, primary PM_{2.5} emissions will be 28% lower than in the baseline projection, and SO₂ emissions decline by 11%. At the same time, greenhouse gas emissions will also be slightly lower (CO₂ by 0.3%, CH₄ by 0.3%) (Table 1).




Fig. 3. PM_{2.5}, SO₂ and NO_x emission reductions in the residential sector in the Household scenario compared with Baseline.

For PM_{2.5}, three groups of measures deliver similar emission reductions: extended natural gas supply in sub-urban areas, clean fuels for cooking in rural areas, and replacement of coal and biomass by briquettes for heating in rural areas. SO₂ reductions also emerge from an increased supply of natural gas to suburban areas, while the introduction of briquettes makes a smaller difference (Fig. 3). All together, these measures will reduce the residential PM_{2.5}, SO₂ and NO_x emissions by 91%, 77% and 47%, respectively, compared with Baseline.

These emission reductions, although they affect only a small share of sources that contribute to ambient PM_{2.5} in the BTH region, will have large benefits for air quality, as they cut low-level emission sources which have a more direct impact on ground-level concentrations and population exposure than emissions from high stacks. Despite being limited to the BTH region, they will reduce mean population exposure by 13% compared to the baseline, i.e., from 68 μg m⁻³ to 59 μg m⁻³. Most importantly, they will allow 81% of people to be exposed to less than 70 μg m⁻³ (i.e., twice the Chinese Ambient Air Quality Standard II and WHO Target Level 1), compared to only 54% of the population in the baseline case (Fig. 4). In addition, they will enable 7.3 million people to switch to clean cooking fuels, so that they will not be exposed to indoor household pollution caused by solid fuel use for cooking any more.

While these measures alone will not be sufficient to meet the air quality standards, they could form an important and effective element of a comprehensive strategy and bring down the contributions from local residential sources to population exposure to about 1 μg m⁻³, from 15 μg m⁻³ in 2010. Four times as much exposure will then

originate from household sources in the provinces surrounding BTH, where similar strategies could deliver comparable benefits. It is noteworthy that there is a strong seasonal component in the emissions from heating, and these sources make a much larger relative contribution during the heating season and especially during high pollution episodes in winter (Liu et al., 2016).

The additional coal use in the power sector resulting from the conservative assumption of maintaining the total demand for natural gas in BTH would increase population exposure by only 0.03 μg m⁻³. Further emission reductions for industry and power plants as well as from agricultural sources in BTH and surrounding areas, are essential to bring exposure levels in compliance with the National Ambient Air Quality standard II.

Beyond the improved compliance with the air quality standards, these emission and exposure reductions will also benefit human health. However, the quantification of health impacts (e.g., number of premature deaths, hospital admissions, absence from work, child mortality) remains uncertain, as most of the available epidemiological evidence is derived from studies in developed countries. To elaborate more robust messages about potential health benefits from policy interventions, we focus on the relative changes in health impacts.

From this, we estimate for the baseline projection that the premature deaths that can be associated with air pollution would decline by 4% between 2010 and 2030. This overall change is a consequence of various factors: (i) total population will increase by 21% in the BTH region; (ii) improved access to clean cooking fuels will reduce mortality




Fig. 4. Proportion of population exposed to different levels of PM_{2.5} concentrations in Household scenario in 2030 (A); source apportionment for population-weighted PM_{2.5} concentrations in the BTH region in the Household scenario in 2030 (B).

from indoor air pollution by 31%; (iii) the mean population exposure to ambient PM_{2.5} will decrease by 20%; (iv) the non-linear shape of the IER curve of the GBD study, which hypothesizes less health improvements from the same absolute reduction of exposure at high concentrations compared to cleaner environments; and (v) the strong aging of the society will make people more susceptible to air pollution, so that premature deaths from ambient pollution are estimated to grow by 32%. In contrast to the rather limited improvement in overall health impacts that can be expected for the baseline scenario, the household scenario would reduce health impacts by almost one-third, with the majority delivered by the elimination of solid fuel use for cooking. In the Household scenario, where the solid fuels used for cooking are completely replaced with LPG and wind power, the premature mortality attributable to household air pollution caused by solid fuel use for cooking is eliminated. This would mostly benefit poorer segments in the society, and especially women and children. Due to the dispersion of PM_{2.5} in the atmosphere, ambient air quality and public health will also improve beyond the BTH area, and we estimate that health benefits outside in the neighboring regions add another 31% to the numbers computed for BTH.

4. Discussion

Faced with the challenge of high air pollution in the densely populated areas, the Chinese government has undertaken significant efforts to reduce emissions from important pollution sources. Early efforts, such as the 11th and 12th Five-Year Plans, focused on SO₂ and NO_x emissions, mainly addressing large point sources in the power and industry sectors, as they are responsible for major shares of total emissions in China. These were complemented by regulations for mobile sources, which are growing rapidly, and small-scale industrial boilers, which relied to a large extent on coal without efficient pollution control equipment. Despite these efforts, episodes of high pollution continued to occur, so that in 2013 the State Council and other authorities issued the *Action Plan* and other regulations, which also included measures for some other sectors.

With a smaller share in energy consumption, pollution from the household and commercial sector received less attention in the past (Liu et al., 2016). However, per unit of energy use, solid fuel combustion in small, low-level sources in the household and commercial sector has significantly stronger impact on ambient air quality and population exposure than large point sources, and its strong seasonality makes it a key source in pollution episodes during the heating season, as confirmed by source apportionment (Yang et al., 2016; Li et al., 2015) and remote sensing (Xiao et al., 2015) studies. Recently, the authorities have begun to address residential emissions from coal use by substituting coal with natural gas and other clean fuels, controlling coal quality and promoting coal briquettes in rural areas. According to the *Action Plan for Comprehensive Management of Air Pollution during Autumn and Winter Season of 2017–2018*, substitution of coal with natural gas and electricity should be implemented in more than 3 million households in the BTH and surrounding regions by 2017 (MEP, 2017).

Previously, we have estimated with the GAINS model that household cooking with solid fuels accounted for 12% of ambient PM_{2.5} concentration globally (Chafe et al., 2014). In this study, focusing on the regional mitigations in Northern China, we build the residential energy use by individual needs in the BTH region, and provide estimates for the potential air quality and human health benefits from implementation of mitigation policies on clean cooking and heating in the residential sector.

A wide range of alternatives is technically available to eliminate the use of solid fuels for cooking and heating in the household and commercial sector, including district heating, grid-based supplies of natural gas for space heating and cooking, LPG for cooking, electricity, biogas, and solar for space heating and cooking. Further expansion of these alternatives depends on policy decisions to prioritize investments in

them. We developed a comprehensive strategy by replacing all the urban solid fuel use with grid-based natural gas, replacing all the rural solid fuel use for cooking with LPG and electricity, and replacing all the rural solid fuel use for space heating with briquettes. Such a re-arrangement of fuel use would substantially reduce the residential PM_{2.5}, SO₂ and NO_x emissions compared with the Baseline (Fig. 3). Although, strictly speaking, coal and biomass briquettes are not clean fuels, they can provide significant reduction of emissions of particulate matter [e.g., Zhi et al., 2008, 2009] and we consider them as short-term to interim alternatives. To meet the space-heating demand in the vast rural areas, the Government suggested regions which do not have proper conditions to use natural gas and electric-power heating devices use coal briquettes and other types of clean fuels as substitutes (MEP, 2017). But further reductions can be achieved when moving to the long-term solutions, such as natural gas, electricity (preferably from renewable sources like wind), and biogas.

Recently, electric-driven heat pumps were proposed as substitutes for household coal-fired heating devices, with preferential policies, such as electricity tariff, purchase subsidy to support the applications (NDRC, 2016), which foresees further reductions beyond our estimation. Xu et al. (2017) estimated that the coal-fired electricity generation and electric-driven heating system can significantly reduce the PM, SO₂ and NO_x emissions by 93%, 97% and 76% respectively, but increase the annualized investment, operation and maintenance cost by \$1.44/m², compared with the direct coal-fired heating system. Therefore, with the implementation of preferential policies and the improvement of technologies, there are large potentials to mitigate air pollution through clean heating and cooking strategies in China.

In addition to the clear benefits of such a strategy for the compliance with ambient air quality standards and the improvement of public health, a rapid adoption of clean heating and cooking fuels will have particular benefits on air quality during the heating season and can significantly reduce the likelihood of extreme pollution episodes during the winter. This is of special relevance for the planning of the 2022 Winter Olympic Games in the BTH area.

Unlike in the earlier events, e.g., the 2008 Olympic Games (Wang et al., 2009), and the 2014 APEC Forum (Tan et al., 2016), heating demand will peak during this time. With large amounts of coal for heating (Zhu et al., 2013) and unfavorable meteorological conditions (Zheng et al., 2015), residential sources contributed about 50% to PM_{2.5} in the urban areas of the BTH region during the January 2013 pollution episode (Li et al., 2015). Our baseline analysis does not foresee much change (a 21% contribution to annual mean exposure in 2030, which approximately doubles during winter episodes). Temporarily restricting emissions from industrial and transport emission sources as successfully mandated for industrial sources and transport in earlier events might not be sufficient to secure acceptable air quality during cold winter periods. Moreover, a temporary shut-down of space heating at the large scale – or of the most polluting space heating sources, i.e., small household devices burning solid fuels, does not seem practical nor socially acceptable. Any solution that does not cause hardship for the population during winter-time must strive for a rapid introduction of clean heating and cooking fuels in a socially acceptable manner. This would also make an important contribution to a long-term strategy towards the PM_{2.5} Interim Targets and Guidelines of the World Health Organization.

Regional collaboration is needed to maximize benefits across the BTH region. Local governments shall evaluate existing plans of gas and electricity supply to residential consumers prioritizing densely populated, highly polluted regions with high reliance on coal. For regions where switching to natural gas and electricity is feasible, supporting funds should be arranged for the construction and upgrading of natural gas and power transmission and distribution networks. Preferential policies such as tax incentives, purchase subsidies can help to decrease the overall costs for the residents. For regions that do not have proper conditions for switching to gas or electricity, coal briquettes and other types of clean fuels can serve as substitutes. Production and supply of

clean coal should be strengthened, and the sales of low quality coal should be prohibited to ensure that coal used in the households meets quality standards. The provincial governments should collaborate with prefectures to evaluate regularly the implementation progress and consider financial support to the less-developed Hebei province.

5. Uncertainties

We applied an integrated approach to highlight the potential opportunities of clean cooking and heating strategies, ranging from energy consumption, future projection, and emission estimation to air quality and human health impacts. Necessarily, the simplifications and assumptions that we had to introduce in order to characterize a complex reality add some uncertainty to quantitative results. Uncertainties emerge from many aspects, including constructing the baseline scenario, the adoption of emission factors, the process of PM_{2.5} concentration modelling, health impact estimation, and the compliance of policies.

We did not perform a quantitative uncertainty analysis, because a comprehensive model to quantify the overall uncertainty of integrated assessment models is not available. Instead, we provided a qualitative analysis of the main sources of uncertainty in the SI. Since we are focused mostly on a scenario comparison, several components of uncertainty can cancel out by the differential analysis. Besides, we use similar emission factors and methodologies with other studies, the magnitude of uncertainty for emissions in GAINS is similar with previous bottom-up emission inventories (Streets et al., 2003; Zhang et al., 2009; Zhao et al., 2013; Li et al., 2017; Wang et al., 2014b). For example, Wang et al. (2014a, 2014b) estimated coefficients of variation for residential sector emissions of SO₂, NO_x, and PM_{2.5} at ± 51%, ± 55%, and ± 68%, respectively, and average 90% confidence interval at about −29% to +45% for SO₂, −31% to +44% for NO_x, and −39% to +49% for PM_{2.5}. So far, the integrated assessment model GAINS has been widely applied in previous studies, including the topics of ammonia mitigation (Zhao et al., 2017) and synthetic natural gas development (Qin et al., 2017) in China.

6. Conclusions

Air pollution is causing a significant health burden in China. Over the last years, policy responses focused on measures to reduce emissions from the power, industry and transportation sectors, with notable success during the 11th and 12th FYP periods. However, we estimate that such strategies will be insufficient to fully reduce ambient PM_{2.5} concentrations below the National Air Quality Standard II of 35 μg m^{−3} throughout the BTH region by 2030, as other emission sources, notably solid fuel combustion for cooking and heating in the household sector, will remain. Furthermore, these sources make large contributions to extreme pollution episodes in winter, which are of great public concern, especially in view of the forthcoming Winter Olympic Games in the BTH area in 2022.

We find that a comprehensive strategy for the residential sector in the BTH region would deliver substantial air quality benefits. Beyond the already planned expansion of district heating and natural gas distribution in urban centers and the foreseen curtailment of coal use for households, such a strategy would

- (i) redirect some of the natural gas that is currently allocated to large industrial facilities towards the residential sector, especially in suburbs with reasonable demand density, thereby eliminating the use of solid fuels and LPG in all households in urban and sub-urban areas (industrial sources might compensate the diverted natural gas by coal),
- (ii) use this LPG, together with additional electricity, to replace biomass cooking in all rural households, and
- (iii) substitute all remaining coal and biomass used for space heating in rural areas by briquettes.

Such a re-arrangement would reduce the residential PM_{2.5}, SO₂ and NO_x emissions by 91%, 77% and 47%, respectively, compared with the Baseline. Although these measures alone will not be sufficient to meet the air quality standards, they could form an important and effective element of a comprehensive mitigation strategy. Similar improvements could be achieved by addressing household sources in the provinces surrounding BTH.

Replacing coal and biomass use for space heating in rural areas with briquettes is a short-term to interim alternative to reduce the present severe air pollution. Further mitigation can be achieved when moving to long-term solutions, such as switching to natural gas, electricity and biogas. Supporting policies and facilities should be well prepared to meet the proper rural heating demand during winter.

Acknowledgement

This work was supported by IIASA's Young Scientists Summer Program (YSSP) sponsored by the National Science Foundation of China (4141101011).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2018.09.059>.

References

- Amann, M., Bertok, I., Borcken, J., Chambers, A., Cofala, J., Dentener, F., et al., 2008. GAINS-Asia. A tool to combat air pollution and climate change simultaneously. Laxenburg, Austria: International Institute for Applied Systems Analysis (IIASA). Available at: <http://pure.iiasa.ac.at/id/eprint/8669/>, Accessed date: 11 April 2018.
- Amann, M., Bertok, I., Borcken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., et al., 2011. Cost-effective control of air quality and greenhouse gases in Europe: modeling and policy applications. *Environ. Model. Softw.* 26 (2), 1489–1501.
- Amann, M., Bertok, I., Borcken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., et al., 2015. Adjusted historic emission data, projections, and optimized emission reduction targets for 2030 - a comparison with COM data 2013. Part A: results for EU-28 - TSAP#16A version 1.1 (no. TSAP report #16A). IIASA. Available at: http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf, Accessed date: 11 April 2018.
- Amann, Markus, Purohit, P., Bhanarkar, A.D., Bertok, I., Borcken-Kleefeld, J., Cofala, J., et al., 2017. Managing future air quality in megacities: a case study for Delhi. *Atmos. Environ.* 161, 99–111. <https://doi.org/10.1016/j.atmosenv.2017.04.041>.
- An, X., Zhu, T., Wang, Z., Li, C., Wang, Y., 2007. A modeling analysis of a heavy air pollution episode occurred in Beijing. *Atmos. Chem. Phys.* 7 (12), 3103–3114.
- Balakrishnan, K., Ghosh, S., Ganguli, B., Sambandam, S., Bruce, N., Barnes, D.F., Smith, K.R., 2013. State and national household concentrations of PM_{2.5} from solid cookfuel use: results from measurements and modelling in India for estimation of the global burden of disease. *Environ. Health* 12 (1), 77. <https://doi.org/10.1186/1476-069X-12-77>.
- Beijing Municipal Government, 2017. The overall plan for Beijing municipality (2016–2035). Available at: <http://zhengwu.beijing.gov.cn/gb/dt/t1494703.htm>, Accessed date: 13 August 2018 (in Chinese).
- Burnett, R.T., Pope III, C.A., Ezzati, M., Olives, C., Lim, S.S., Mehta, S., et al., 2014. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ. Health Perspect.* <https://doi.org/10.1289/ehp.1307049>.
- Cao, G., Zhang, X., Wang, Y., Zheng, F., 2008. Estimation of emissions from field burning of crop straw in China. *Chin. Sci. Bull.* 53, 784–790. <https://doi.org/10.1007/s11434-008-0145-4>.
- Chafe, Z.A., Brauer, M., Klimont, Z., Van Dingenen, R., Mehta, S., Rao, S., Riahi, K., Dentener, F., Smith, K.R., 2014. Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease. *Environ. Health Perspect.* 122, 1314–1320. <https://doi.org/10.1289/ehp.1206340>.
- Chen, Y., Zhi, G., Feng, Y., Liu, D., Zhang, G., Li, J., Sheng, G., Fu, J., 2009. Measurements of black and organic carbon emission factors for household coal combustion in China: implication for emission reduction. *Environ. Sci. Technol.* 43, 9495–9500. <https://doi.org/10.1021/es9021766>.
- Chen, Z., Wang, J.N., Ma, G.X., Zhang, Y.S., 2013. China tackles the health effects of air pollution. *Lancet* 382 (9909), 1959–1960.
- China Electric Power Yearbook Editorial Committee, 2011. *China Electric Power Yearbook*. China Electric Power Press, Beijing.
- China Electric Power Yearbook Editorial Committee, 2015. *China Electric Power Yearbook*. China Electric Power Press, Beijing.
- Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., et al., 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389 (10082), 1907–1918. [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6).

- DHUDH (Department of Housing and Urban-rural Development of Hebei Province), 2017. Hebei Province urban system plan. Available at: http://www.hebjs.gov.cn/zhengcewenjian/tingfawenjian/201703/t20170307_226738.html, Accessed date: 13 August 2018 (in Chinese).
- Forouzanfar, M., et al., 2015. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 386 (10010), 2287–2323.
- Geng, G., Zhang, Q., Tong, D., Li, M., Zheng, Y., Wang, S., He, K., 2017. Chemical composition of ambient PM_{2.5} over China and relationship to precursor emissions during 2005–2012. *Atmos. Chem. Phys.* 17 (14), 9187–9203. <https://doi.org/10.5194/acp-17-9187-2017>.
- HPDHURC (Hebei Provincial Department of Housing and Urban Rural Construction), 2017. The 13th five-year plan for Hebei urban heating. Available at: http://www.hebjs.gov.cn/xgzz/cjc/tfwj_1_1001/201705/t20170526_228033.html, Accessed date: 13 August 2018 (in Chinese).
- International Energy Agency (IEA), 2015. *World Energy Outlook 2015*. Paris, France.
- International Energy Agency (IEA), 2016. *World energy outlook special report 2016: energy and air pollution*. Available at: <http://www.iea.org/publications/freepublications/publication/weo-2016-special-report-energy-and-air-pollution.html>, Accessed date: 13 August 2018.
- Kiesewetter, G., Borken-Kleefeld, J., Schöpp, W., Heyes, C., Thunis, P., Bessagnet, B., et al., 2015. Modelling street level PM₁₀ concentrations across Europe: source apportionment and possible futures. *Atmos. Chem. Phys.* 15 (3), 1539–1553. <https://doi.org/10.5194/acp-15-1539-2015>.
- Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M., 2009. Projections of SO₂, NO_x and carbonaceous aerosols emissions in Asia. *Tellus Ser. B Chem. Phys. Meteorol.* 61, 602–617. <https://doi.org/10.1111/j.1600-0889.2009.00428.x>.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., Schöpp, W., 2017. Global anthropogenic emissions of particulate matter including black carbon. *Atmos. Chem. Phys.* 17 (14), 8681–8723.
- Li, X., Wang, S., Duan, L., Hao, J., Nie, Y., 2009. Carbonaceous aerosol emissions from household biofuel combustion in China. *Environ. Sci. Technol.* 43, 6076–6081. <https://doi.org/10.1021/es803330j>.
- Li, X., Zhang, Q., Zhang, Y., Zheng, B., Wang, K., Chen, Y., et al., 2015. Source contributions of urban PM_{2.5} in the Beijing–Tianjin–Hebei region: changes between 2006 and 2013 and relative impacts of emissions and meteorology. *Atmos. Environ.* 123 (Part A), 229–239. <https://doi.org/10.1016/j.atmosenv.2015.10.048>.
- Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D.G., Carmichael, G.R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., Zheng, B., 2017. MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP. *Atmos. Chem. Phys.* 17, 935–963. <https://doi.org/10.5194/acp-17-935-2017>.
- Liang, L., Wu, W., Lal, R., Guo, Y., 2013. Structural change and carbon emission of rural household energy consumption in Huanai, northern China. *Renew. Sust. Energ. Rev.* 28, 767–776. <https://doi.org/10.1016/j.rser.2013.07.041>.
- Liu, F., Klimont, Z., Zhang, Q., Cofala, J., Zhao, L., Huo, H., et al., 2013. Integrating mitigation of air pollutants and greenhouse gases in Chinese cities: development of GAINS-City model for Beijing. *J. Clean. Prod.* <https://doi.org/10.1016/j.jclepro.2013.03.024>.
- Liu, J., Mauzerall, D.L., Chen, Q., Zhang, Q., Song, Y., Peng, W., et al., 2016. Air pollutant emissions from Chinese households: a major and underappreciated ambient pollution source. *Proc. Natl. Acad. Sci.* 113 (28), 7756–7761.
- Lutz, W., Butz, W.P., S., K.C., 2014. *World Population & Human Capital in the Twenty-first Century*. Oxford University Press, UK.
- Ma, Q., Cai, S., Wang, S., Zhao, B., Martin, R.V., Brauer, M., et al., 2017. Impacts of coal burning on ambient PM_{2.5} pollution in China. *Atmos. Chem. Phys.* 17 (7), 4477–4491. <https://doi.org/10.5194/acp-17-4477-2017>.
- Ministry of Environmental Protection of China (MEP), 2012. The twelfth five-year plan on air pollution prevention and control in key areas. Available at: http://www.mep.gov.cn/gkml/hbb/bwj/201212/t20121205_243271.htm, Accessed date: 13 August 2018 (in Chinese).
- Ministry of Environmental Protection of China (MEP), 2013. The state council issues action plan on prevention and control of air pollution introducing ten measures to improve air quality. Available at: http://english.mep.gov.cn/News_service/infocus/201309/t20130924_260707.htm, Accessed date: 13 August 2018 (in Chinese).
- Ministry of Environmental Protection of China (MEP), 2017. Comprehensive management of air pollution during autumn and winter season of 2017–2018. Available at: http://www.zhb.gov.cn/gkml/hbb/bwj/201708/t20170824_420330.htm, Accessed date: 13 August 2018 (in Chinese).
- National Bureau of Statistics, 2011. *China Energy Statistical Yearbook*. China Statistics Press, Beijing.
- National Bureau of Statistics, 2016. *China Statistical Yearbook*. China Statistics Press, Beijing.
- NDRC (National Development and Reform Commission), 2016. Promoting the alternative of electric power's guidance. Available at: http://www.gov.cn/xinwen/2016-05/25/content_5076579.htm, Accessed date: 17 July 2017 (In Chinese).
- Qin, Y., Wagner, F., Scovronick, N., Peng, W., Yang, J., Zhu, T., Smith, K.R., Mauzerall, D.L., 2017. Air quality, health, and climate implications of China's synthetic natural gas development. *PNAS* 114, 4887–4892. <https://doi.org/10.1073/pnas.1703167114>.
- Ru, M., Tao, S., Smith, K., Shen, G., Shen, H., Huang, Y., et al., 2015. Direct energy consumption associated emissions by rural-to-urban migrants in Beijing. *Environ. Sci. Technol.* 49 (22), 13708–13715. <https://doi.org/10.1021/acs.est.5b03374>.
- Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L.D., Fagerli, H., et al., 2012. The EMEP MSC-W chemical transport model—technical description. *Atmos. Chem. Phys.* 12, 7825–7865.
- Streets, D.G., Bond, T.C., Carmichael, G.R., Fernandes, S.D., Fu, Q., He, D., Klimont, Z., Nelson, S.M., Tsai, N.Y., Wang, M.Q., Woo, J.H., Yarber, K.F., 2003. An inventory of gaseous and primary aerosol emissions in Asia in the year 2000. *J. Geophys. Res.-Atmos.* 108, 8809. <https://doi.org/10.1029/2002JD003093>.
- Tan, Z., Wang, Y., Ye, C., Zhu, Y., Li, Y., Liang, P., et al., 2016. Evaluating vehicle emission control policies using on-road mobile measurements and continuous wa-velet transform: a case study during the Asia-Pacific economic cooperation forum, China 2014. *Atmos. Chem. Phys. Discuss.* 1–39. <https://doi.org/10.5194/acp-2016-460>.
- Tianjin Municipal Government, 2006. The overall plan for Beijing municipality (2005–2020). Available at: http://www.tj.gov.cn/xw/tztg/200610/t20061022_270146.html, Accessed date: 13 August 2018 (in Chinese).
- Wang, M., Zhu, T., Zheng, J., Zhang, R.Y., Zhang, S.Q., Xie, X.X., et al., 2009. Use of a mobile laboratory to evaluate changes in on-road air pollutants during the Beijing 2008 Summer Olympics. *Atmos. Chem. Phys.* 9 (21), 8247–8263. <https://doi.org/10.5194/acp-9-8247-2009>.
- Wang, J., Lei, Y., Yang, J., Yan, G., 2012. China's air pollution control calls for sustainable strategy for the use of coal. *Environ. Sci. Technol.* 46, 4263–4264. <https://doi.org/10.1021/es301226n>.
- Wang, Y., Ying, Q., Hu, J., Zhang, H., 2014a. Spatial and temporal variations of six criteria air pollutants in 31 provincial capital cities in China during 2013–2014. *Environ. Int.* 73, 413–422. <https://doi.org/10.1016/j.envint.2014.08.016>.
- Wang, S.X., Zhao, B., Cai, S.Y., Klimont, Z., Nielsen, C.P., Morikawa, T., Woo, J.H., Kim, Y., Fu, X., Xu, J.Y., Hao, J.M., He, K.B., 2014b. Emission trends and mitigation options for air pollutants in East Asia. *Atmos. Chem. Phys.* 14, 6571–6603. <https://doi.org/10.5194/acp-14-6571-2014>.
- World Bank, 2007. *Cost of pollution in China: economic estimates of physical damages*. Available at: <http://documents.worldbank.org/curated/en/782171468027560055/Cost-of-pollution-in-China-economic-estimates-of-physical-damages>, Accessed date: 11 April 2018.
- World Bank, 2016. *The Cost of Air Pollution: Strengthening the Economic Case for Action* (No. 108141). pp. 1–122. Available at: <http://documents.worldbank.org/curated/en/781521473177013155/The-cost-of-air-pollution-strengthening-the-economic-case-for-action> (Accessed on 11 April 2018).
- World Health Organization (WHO), 2016. *Ambient air pollution: a global assessment of exposure and burden of disease*. Geneva, Switzerland. Available at: <http://www.who.int/phe/publications/air-pollution-global-assessment/en/>, Accessed date: 11 April 2018.
- Xiao, Q., Ma, Z., Li, S., Liu, Y., 2015. The impact of winter heating on air pollution in China. *PLoS ONE* 10 (1), e0117311. <https://doi.org/10.1371/journal.pone.0117311>.
- Xu, C., Wang, C., Xu, G., Hu, Y., Guo, H., Yang, Y., 2017. Thermodynamic and environmental evaluation of an improved heating system using electric-driven heat pumps: a case study for Jing-Jin-Ji region in China. *J. Clean. Prod.* 165, 36–47. <https://doi.org/10.1016/j.jclepro.2017.07.087>.
- Yang, H., Chen, J., Wen, J., Tian, H., Liu, X., 2016. Composition and sources of PM_{2.5} around the heating periods of 2013 and 2014 in Beijing: implications for efficient mitigation measures. *Atmos. Environ.* 124 (Part B), 378–386. <https://doi.org/10.1016/j.atmosenv.2015.05.015>.
- Zhang, Q., Streets, D.G., Carmichael, G.R., He, K.B., Huo, H., Kannari, A., Klimont, Z., Park, I.S., Reddy, S., Fu, J.S., Chen, D., Duan, L., Lei, Y., Wang, L.T., Yao, Z.L., 2009. Asian emissions in 2006 for the NASA INTEX-B mission. *Atmos. Chem. Phys.* 9, 5131–5153.
- Zhang, Q., He, K., Huo, H., 2012. Policy: cleaning China's air. *Nature* 484 (7393), 161–162. <https://doi.org/10.1038/484161a>.
- Zhao, B., Wang, S.X., Liu, H., Xu, J.Y., Fu, K., Klimont, Z., Hao, J.M., He, K.B., Cofala, J., Amann, M., 2013. NO_x emissions in China: historical trends and future perspectives. *Atmos. Chem. Phys.* 13, 9869–9897. <https://doi.org/10.5194/acp-13-9869-2013>.
- Zhao, Z.Q., Bai, Z.H., Winiwarter, W., Kiesewetter, G., Heyes, C., Ma, L., 2017. Mitigating ammonia emission from agriculture reduces PM_{2.5} pollution in the Hai River Basin in China. *Sci. Total Environ.* 609, 1152–1160. <https://doi.org/10.1016/j.scitotenv.2017.07.240>.
- Zheng, X., Wei, C., Qin, P., Guo, J., Yu, Y., Song, F., Chen, Z., 2014. Characteristics of residential energy consumption in China: findings from a household survey. *Energy Policy* 75 (0), 126–135. <https://doi.org/10.1016/j.enpol.2014.07.016>.
- Zheng, G.J., Duan, F.K., Su, H., Ma, Y.L., Cheng, Y., Zheng, B., et al., 2015. Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous reactions. *Atmos. Chem. Phys.* 15 (6), 2969–2983. <https://doi.org/10.5194/acp-15-2969-2015>.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., et al., 2018. Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions. *Atmos. Chem. Phys. Discuss.* 2018, 1–27. <https://doi.org/10.5194/acp-2018-374>.
- Zhi, G., Chen, Y., Feng, Y., Xiong, S., Li, J., Zhang, G., Sheng, G., Fu, J., 2008. Emission characteristics of carbonaceous particles from various residential coal-stoves in China. *Environ. Sci. Technol.* 42, 3310–3315. <https://doi.org/10.1021/es702247q>.
- Zhi, G., Peng, C., Chen, Y., Liu, D., Sheng, G., Fu, J., 2009. Deployment of coal briquettes and improved stoves: possibly an option for both environment and climate. *Environ. Sci. Technol.* 43, 5586–5591. <https://doi.org/10.1021/es802955d>.
- Zhu, D., Tao, S., Wang, R., Shen, H., Huang, Y., Shen, G., et al., 2013. Temporal and spatial trends of residential energy consumption and air pollutant emissions in China. *Appl. Energy* 106, 17–24. <https://doi.org/10.1016/j.apenergy.2013.01.040>.