

1 **The generalization of Gulland's method: how to estimate maturity ogives**  
2 **when juvenile data are missing while spawner demography is known**

3 Mikko Heino<sup>a,b,c,d,\*</sup>, Olav Rune Godø<sup>b,e</sup>, Ulf Dieckmann<sup>c</sup>

4 <sup>a</sup>Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen, Nor-  
5 way

6 <sup>b</sup>Institute of Marine Research, PO Box 1879 Nordnes, N-5817 Bergen, Norway

7 <sup>c</sup>Evolution and Ecology Program, International Institute for Applied Systems Analysis (IIASA),  
8 A-2361 Laxenburg, Austria

9 <sup>d</sup>Institute of Oceanography, National Taiwan University, Taipei 106, Taiwan

10 <sup>e</sup>NORCE Norwegian Research Centre AS, PO Box 22 Nygårdstangen, N-5838 Bergen, Nor-  
11 way

12 \*Corresponding author: tel +47 5558 4544, fax +47 5558 4450, email [mikko.heino@uib.no](mailto:mikko.heino@uib.no)

13

## 14 **Abstract**

15 The proportions of mature individuals at age or length, collectively known as the maturity ogive,  
16 are a key population characteristic and serve as critical input to age-disaggregated stock assess-  
17 ments. John Gulland showed in 1964 that it is possible to estimate maturity ogives even when  
18 representative data on immature individuals are not available, provided that one can distinguish  
19 newly mature individuals (first-time spawners) from those that had matured earlier (repeat  
20 spawners). Gulland's method offers a valuable tool for obtaining information on an unobserved  
21 part of a population and is also applicable to other ontogenetic transitions, such as metamor-  
22 phosis, smolting, ontogenetic niche shifts, and sex change. Here we present a full derivation of  
23 Gulland's method from first principles, applicable to the general case in which the survival of  
24 immature, first-spawning, and repeat-spawning individuals may differ. Better observation  
25 methods, in particular in sclerochronology and histology, are expected to make meeting this  
26 method's data requirements—i.e., the separation of first-time and repeat spawners—more often  
27 achievable, and estimating maturity ogives could serve as an additional incentive for allocating  
28 resources to enhanced data collection. With the generalization presented here, we hope to make  
29 Gulland's method better known and more widely accessible.

30

31 **Keywords:** maturity ogives, life-history transitions, reproductive potential, stock dynamics

## 32 **1. Introduction**

33 Maturity ogives measure the proportions of mature individuals at age or length and serve as a  
34 key characteristic of populations, directly impacting their reproductive potential (Murua and  
35 Saborido-Rey, 2003; ICES, 2008; Flores et al., 2015). The demographic structure of the mature  
36 part of a population has profound consequences for recruitment and population dynamics  
37 (Trippel et al., 1997; Ottersen et al., 2006; Köster et al., 2013; Hixon et al., 2014). At the same

38 time, fish maturation is a highly plastic process that is influenced by an array of environmental  
39 factors such as food availability and temperature (Stearns and Crandall, 1984; Trippel, 1995).  
40 This implies that it is important to see maturity ogives as dynamic, rather than static, population  
41 characteristics and that an accurate understanding of stock dynamics necessitates the regular  
42 updating of a stock's maturity ogive.

43         When representative measurements of both immature and mature individuals are avail-  
44 able, it is straightforward to estimate a population's maturity ogive as the proportions of mature  
45 individuals among all individuals, mature and immature, across all age or length classes. Indeed,  
46 it would seem obvious that data on both immature and mature individuals were always needed  
47 for estimating maturity ogives. However, John Gulland has shown, already more than 50 years  
48 ago, that this is not necessary: age-dependent maturity ogives can be calculated based on age-  
49 specific proportions of first-time spawning individuals among all spawning individuals (Gul-  
50 land, 1964). This is potentially a very important methodological discovery, because obtaining  
51 representative samples of both immature and mature individuals can be difficult. A number of  
52 challenges are evident: immature and mature fish may be spatially segregated, sampling gear  
53 may have reduced catchability for smaller sizes, and—when obtaining data from commercial  
54 fisheries—fishermen often face regulations specifically designed to reduce catching juvenile  
55 fish that may end up being discarded. Gulland's method therefore offers considerable promise  
56 in the many situations in which only the mature part of a population is amenable to quantitative  
57 sampling.

58         In reality, Gulland's method has seen only sporadic use, probably because it is rare that  
59 first-time and repeat spawners are separated as part of routine stock monitoring. However, in  
60 some fish populations, the age at first spawning can be estimated from scales or otoliths based  
61 on so-called spawning checks. This is the case for Northeast Arctic cod (*Gadus morhua*), for  
62 which spawning checks can be identified in otoliths (Rollefsen, 1933; Zuykova et al., 2009).

63 Indeed, Northeast Arctic cod is the stock for which Gulland first devised his method. Since then,  
64 the method has recurrently been applied to this commercially and economically important stock  
65 (Jørgensen, 1990; Heino et al., 2002; Svåsand et al., 2003; Zuykova et al., 2009; Yaragina, 2010)  
66 and contributes to the maturity ogives used in the official stock assessment (Zuykova et al.,  
67 2009; ICES, 2017). Another important example of such applications is Norwegian spring-  
68 spawning herring (*Clupea harengus*), for which scales can be used to identify first-time spawn-  
69 ers (Lea, 1928; Runnström, 1936; Engelhard et al., 2003). Gulland's method has been applied  
70 to this stock in a few publications (Engelhard and Heino, 2004a, 2004b) and, since 2010, in the  
71 official stock assessment (ICES, 2016). Species for which Gulland's method has not yet been  
72 used, even though first spawning can be identified from otoliths or scales, are as diverse as  
73 shortnose sturgeon (*Acipenser brevirostrum*; Taubert, 1980), steelhead trout (*Oncorhynchus*  
74 *mykiss*; Narver, 1969), Atlantic halibut (*Hippoglossus hippoglossus*; Devold, 1938), and orange  
75 roughy (*Hoplostethus atlanticus*; Francis and Horn, 1997). Future developments in sclerochro-  
76 nology could make this list much longer.

77 Gulland (1964) presented his method through a worked example rather than in terms of  
78 general equation(s). He also noted that differential survival between immature and mature fish  
79 will bias the results, but considered this bias unimportant and did not present a correction. Terje  
80 Jørgensen (1990) was the first to express Gulland's method as a general equation. He also pre-  
81 sented a way to account for the difference in survival between mature and immature fish, albeit  
82 indirectly, by adjusting the counts of repeat spawners. However, he did not present the deriva-  
83 tion of the equations, nor did he allow for the survival of first-time spawners to differ from the  
84 survival of repeat spawners. Here we present a full derivation of Gulland's method from first  
85 principles, applicable to the general case in which the survival of immature, first-spawning, and  
86 repeat-spawning individuals may differ. With this generalization, we hope to make Gulland's  
87 method better known and more widely accessible.

## 88 2. Logic of Gulland's method

89 We first describe the general idea underlying Gulland's method. When survival of all types of  
90 individuals of the same age is equal, the method is straightforward and can easily be illustrated  
91 graphically (Fig. 1). This simplifying assumption is then relaxed in the next section.

92 The goal of Gulland's method is to estimate the age-specific proportions  $o_t$  of mature  
93 individuals for the population's cohorts at each age  $t$  from the observed (sampled) numbers of  
94 first-spawning and repeat-spawning fish,  $n_t^F$  and  $n_t^R$ , respectively. Since we are dealing with  
95 each cohort separately, the index  $t$  can be interpreted as either age or time (both measured in  
96 numbers of spawning seasons), whichever is more convenient. An individual is born as a juve-  
97 nile (life stage J), then matures and becomes a first-time spawner (F), before turning into a  
98 repeat spawner (R) for the rest of its life. Thus, the generalized ontogeny is of type  $J \rightarrow \dots \rightarrow$   
99  $J \rightarrow F \rightarrow R \rightarrow \dots \rightarrow R \rightarrow \dagger$ , with each arrow corresponding to a time step, typically one year. As  
100 appropriate, other conventions for naming the successive stages can be adopted, as long as the  
101 distinction between juveniles, newly mature individuals, and individuals that were newly ma-  
102 ture during earlier observation steps is maintained.

103 Gulland's method is iterative, progressing backward in time. The estimation is started  
104 from the earliest age at which a cohort is fully mature. In the absence of data on juveniles, this  
105 can be identified as the latest age at which the samples contain first-time spawners. We there-  
106 fore take this as the reference time and denote it by  $T$  (Fig. 1). At time  $T$ , the maturity ogive,  
107 by definition, has the value  $o_T = 1$ . We can then work backward in time by noting that the  
108 proportion  $o_{T-1}$  of mature individuals among all individuals one time step earlier, i.e., at time  
109  $T - 1$ , equals the proportion  $r_T$  of repeat spawners among all mature individuals at time  $T$ ,  
110  $o_{T-1} = r_T$ . The latter proportion is known directly from the sampling,  $r_T = n_T^R / (n_T^F + n_T^R)$ . For  
111 time  $T - 2$ , the proportion  $o_{T-2}$  of mature individuals among all individuals likewise equals  
112 the proportion of repeat spawners among all individuals at time  $T - 1$  (Fig. 1). This proportion

113 is not known from the outset. However, the proportion  $o_{T-1}$  of mature individuals among all  
 114 individuals in the cohort is known from the previous step, and the proportion  $r_{T-1}$  of repeat  
 115 spawners among all mature individuals is again known from the sampling. We can multiply  
 116 these two proportions to determine  $o_{T-2} = o_{T-1}r_{T-1}$  (Fig. 1). This equation is readily general-  
 117 ized for any time  $t$ . Thus, we obtain the general iterative equation for Gulland's method under  
 118 uniform survival probabilities,

$$o_{t-1} = o_t r_t, \quad (1a)$$

119 or equivalently,  $o_t = o_{t+1} r_{t+1}$ . Accumulating all iterations, the maturity ogive can be expressed  
 120 as an explicit closed-form function of time  $t$ ,

$$o_t = \prod_{\tau=t+1}^T r_{\tau}. \quad (1b)$$

### 121 3. Generalization of Gulland's method

122 We now derive Gulland's method from first principles. This allows considering the role of  
 123 survival, and of differences in survival between life stages, in a rigorous way.

124 Changes in the abundances of fish in a cohort over one time step are described as follows,

$$N_{t+1}^J = s_t^{JJ} N_t^J (1 - m_t), \quad (2a)$$

$$N_{t+1}^F = s_t^{JF} N_t^J m_t, \quad (2b)$$

$$N_{t+1}^R = s_t^{FR} N_t^F + s_t^{RR} N_t^R, \quad (2c)$$

125 where  $N_t^J$ ,  $N_t^F$ , and  $N_t^R$  are the abundances of juvenile, first-spawning, and repeat-spawning fish  
 126 at time  $t$ ,  $s_t^{JJ}$ ,  $s_t^{JF}$ ,  $s_t^{FR}$ , and  $s_t^{RR}$  are the survival probabilities for the transitions  $J \rightarrow J$ ,  $J \rightarrow F$ ,  
 127  $F \rightarrow R$ ,  $R \rightarrow R$ , respectively, and  $m_t$  is the probability of maturation during the next time step.  
 128 Here we have assumed that, at any time  $t$ , the survival probability for transitions  $R \rightarrow R$  is in-  
 129 dependent of the number of preceding spawning events.

130 The maturity ogive can always be expressed in terms of stage-specific abundances:

$$o_t = \frac{N_t^F + N_t^R}{N_t^J + N_t^F + N_t^R} = \frac{1}{1 + \frac{N_t^J}{N_t^F + N_t^R}}. \quad (3)$$

131 With expressions derived from Equations 2a–2c, the terms  $N_t^J$  and  $N_t^F + N_t^R$  can be expressed

132 in terms of the cohort composition at time  $t + 1$ . First, rearranging Equation 2b gives  $N_t^J =$

133  $N_{t+1}^F / s_t^{JF} / m_t$ . The maturation probability  $m_t$  can be solved from Equations 2a and 2b as

$$m_t = \frac{1}{1 + \frac{s_t^{JF} N_{t+1}^J}{s_t^{JJ} N_{t+1}^F}}. \quad (4a)$$

134 Substituting Equation 4a into the expression for  $N_t^J$  gives

$$N_t^J = \frac{N_{t+1}^F \left(1 + \frac{s_t^{JF} N_{t+1}^J}{s_t^{JJ} N_{t+1}^F}\right)}{s_t^{JF}}. \quad (4b)$$

135 Second, dividing Equation 2c with  $N_t^F + N_t^R$ , using the definition  $r_t = N_t^R / (N_t^F + N_t^R)$ , and

136 rearranging gives

$$N_t^F + N_t^R = \frac{N_{t+1}^R}{s_t^{FR}(1-r_{t+1}) + s_t^{RR}r_{t+1}}. \quad (4c)$$

137 Inserting Equations 4b and 4c into Equation 3 gives

$$o_t = \frac{1}{1 + \frac{N_{t+1}^F}{N_{t+1}^R} \left(1 + \frac{s_t^{JF} N_{t+1}^J}{s_t^{JJ} N_{t+1}^F}\right) \frac{s_t^{FR}(1-r_{t+1}) + s_t^{RR}r_{t+1}}{s_t^{JF}}}. \quad (5a)$$

138 This equation still contains two unknown ratios of abundances. As ratios, they are independent

139 of total abundance and can instead be expressed solely in terms of parameters  $o_{t+1}$  and  $r_{t+1}$ .

140 Specifically,  $N_{t+1}^F / N_{t+1}^R = o_{t+1}(1 - r_{t+1}) / o_{t+1}r_{t+1}$  and

141  $N_{t+1}^J / N_{t+1}^F = (1 - o_{t+1}) / o_{t+1}(1 - r_{t+1})$ . Using these relationships, rearranging, and shifting

142 the time index back by one time step gives

$$o_{t-1} = \frac{o_t r_t}{o_t r_t + \left( o_t (1 - r_t) + (1 - o_t) \frac{s_{t-1}^{JF}}{s_{t-1}^{JJ}} \right) \left( \frac{s_{t-1}^{FR}}{s_{t-1}^{JF}} (1 - r_t) + \frac{s_{t-1}^{RR}}{s_{t-1}^{JF}} r_t \right)}. \quad (5b)$$

143 This iterative equation allows determining the value  $o_{t-1}$  of the maturity ogive at time  $t - 1$   
 144 from three sources of information: (i) the ratio  $r_t$  of repeat spawners among all mature individ-  
 145 uals, known from (representative) sampling, (ii) three ratios of survival probabilities, possibly  
 146 known from independent observations, and (iii) the value  $o_t$  of the maturity ogive at time  $t$ ,  
 147 known from the equation's previous iteration or because the cohort is known to be fully mature  
 148 at time  $t$ . Equation 5b is similar to Equation 1a, but includes a correction in form of the denom-  
 149 inator, which accounts for differences in survival between the life stages. It is readily seen that  
 150 Equation 5b reduces to Equation 1a as a special case when all survival probabilities are equal.

151 Equation 5b contains three ratios of survival probabilities that all include  $s_{t-1}^{JF}$ , which  
 152 therefore naturally serves as the reference against which the other survival probabilities can be  
 153 compared. Sometimes it is more convenient to use another survival probability as the reference:  
 154 for instance, when we use  $s_{t-1}^{RR}$  as the reference, Equation 5b becomes

$$o_{t-1} = \frac{o_t r_t}{o_t r_t + \left( o_t (1 - r_t) \frac{s_{t-1}^{RR}}{s_{t-1}^{JF}} + (1 - o_t) \frac{s_{t-1}^{RR}}{s_{t-1}^{JJ}} \right) \left( \frac{s_{t-1}^{FR}}{s_{t-1}^{RR}} (1 - r_t) + r_t \right)}. \quad (5c)$$

155 Figure 2 illustrates the sensitivity of the estimated maturity ogive to departures from  
 156 equal age-specific survival between maturity stages. Not surprisingly, the sensitivity is greater  
 157 when the estimated maturity proportion is near the middle of the possible range (Fig. 2, left  
 158 panels) compared to when the proportion is near the border of the possible range (Fig. 2, right  
 159 panels). The general tendency is that assuming equal survival ratios between maturity stages  
 160 leads to positively biased maturity estimates (red colors in Fig. 2) when juvenile survival is low  
 161 relative to repeat-spawning survival ( $s_{t-1}^{JJ}/s_{t-1}^{RR} < 1$ ) and/or when juvenile-to-first-spawning  
 162 survival is low relative to repeat-spawning survival ( $s_{t-1}^{JF}/s_{t-1}^{RR} < 1$ ); the opposite is true when



163 these ratios are high.

#### 164 **4. Example: Northeast Arctic cod**

165 We illustrate the generalized Gulland's method for the 1928 cohort of Northeast Arctic cod, the  
166 first cohort for which suitable data are available. For this stock, historic samples are available  
167 from the spawning grounds, distinguishing first-time spawners and repeat spawners, but no  
168 representative data are available for juvenile individuals (e.g., Jørgensen, 1990; Heino et al.,  
169 2002). For the 1928 cohort, the proportions of repeat spawners among all mature individuals  
170 equals 0 for ages 6 years and younger, equals 0.03, 0.24, 0.47, 0.66, 0.70, 0.85, 0.96, and 0.95  
171 for ages 7 to 14 years, respectively, and equals 1 for older individuals. Because only mature  
172 fish were subject to fishing in the spawning grounds, it is likely that juveniles experienced a  
173 higher age-specific survival ( $s_t^{JJ}$ ) than fish that entered the spawning grounds ( $s_t^{JF}$ ,  $s_t^{FR}$ , and  
174  $s_t^{RR}$ ). We therefore let the ratio  $s_t^{JJ}/s_t^{RR}$  vary. In addition to considering the case  $s_t^{JF}/s_t^{RR} = 1$   
175 and  $s_t^{FR}/s_t^{RR} = 1$ , we examine a scenario in which individuals lacking spawning experience  
176 suffer from additional mortality during their spawning migration, by considering the case  
177  $s_t^{JF}/s_t^{RR} = 0.7$  and  $s_t^{FR}/s_t^{RR} = 0.9$ ; the particular numbers here are chosen for illustrative pur-  
178 poses only.

179 Applying our generalization of Gulland's method reveals that the estimated maturity  
180 ogive is potentially sensitive to departures from equal age-specific survival between maturity  
181 stages (Fig. 2, 3). For instance, when spawning individuals suffer from higher mortality than  
182 those that do not spawn, assuming that no such survival difference exists results in an ogive  
183 that is downward biased. If the survival difference is large ( $s_t^{JJ}/s_t^{RR} = 2$ ), the true ogive is up  
184 to about 15 percent points higher than the estimated one (Fig. 3a). If we assume that lack of  
185 spawning experience reduces survival, the maximal bias when  $s_t^{JJ}/s_t^{RR} = 2$  is slightly lower, at

186 about ten percent points (Fig. 3b).

187 A potential source of bias for estimating the maturity ogive of Northeast Arctic cod is  
188 skipped spawning (Jørgensen et al., 2006; Skjæraasen et al., 2012). The effect of skipped  
189 spawning on spawner demography is that the sampled proportions of repeat spawners among  
190 all mature individuals ( $r_T$ ) are less than their true proportions. The strength of this bias will vary  
191 with a cohort's age, because skipped spawning depends on spawning experience. Figure 4a  
192 shows that if skipped spawning is very frequent and leads to a serious underrepresentation of  
193 repeat spawners, the estimated maturity ogive can be seriously biased downward. However, for  
194 the documented levels of skipped spawning (~24 % in 2006–2008, Skjæraasen et al., 2012), the  
195 bias is modest, at most seven percent points (Fig. 4a for  $x = 0.25$ ).

196 Another possible source of uncertainty is the misidentification of first-time and repeat  
197 spawners. Figure 4b shows that misidentifying first-time spawners as second-time spawners or  
198 vice versa at a relatively high rate (20%) results in a modest downward bias in the estimated  
199 maturity ogive. The largest error is about seven percent points for ages 10–11 years.

## 200 **5. Discussion**

201 Here we have presented a derivation of the generalized Gulland's method to estimate maturity  
202 ogives in the absence of data on juveniles. The information required instead are age-specific  
203 proportions of repeat spawners among spawning individuals, as well as ratios of age-specific  
204 survival among juveniles, maturing individuals, first-time spawners, and repeat spawners, all  
205 for a given cohort. These survival ratios can be based on independent observations or expert  
206 knowledge. This is more practical than the correction proposed by Jørgensen (1990), which  
207 requires adjusting the input data before applying the original Gulland's method assuming stage-  
208 independent survival (his Equation 3 and our Equation 1a).

209 As underscored by our examples above, assuming the aforementioned survival ratios to

210 equal 1 can greatly bias the estimation of maturity ogives when these ratios in fact significantly  
211 differ from 1. While estimating survival in wild populations is always difficult, two mitigating  
212 considerations are worth emphasizing. First, it is only the aforementioned survival ratios that  
213 enter the generalized Gulland's method, not the absolute values of survival probabilities. Sec-  
214 ond, these ratios will deviate from 1 only when maturation stages differentially impact age-  
215 specific survival. When such effects of maturation stages within each age class are weak, as-  
216 suming the ratios to equal 1 will not cause major biases in the estimated maturity ogives.

217         Because maturity ogives result from maturation dynamics of a cohort of individuals, the  
218 most natural biological unit for estimating ogives is a cohort. For this reason, we have presented  
219 the derivation here for cohorts, as did Gulland (1964) and Jørgensen (1990). However, Gul-  
220 land's method—and estimation of maturity ogives in general—can also be used with data from  
221 a single year or with data averaged over a range of years. The implicit assumption is that age-  
222 specific changes in maturity observed for concurrently recorded age groups is similar to what  
223 would have been observed when following a cohort over time. The situation is analogous to  
224 growth curves, which can be estimated either by cohort or by year (Beverton and Holt, 1957, p.  
225 282; Gulland, 1969, p. 93; Ricker, 1975, p. 205). The disadvantage of any such estimations by  
226 year is that factors that are specific to a cohort will confound the detection of age-specific  
227 change, be it in maturity or in size. For example, strong year classes of Norwegian spring-  
228 spawning herring show different maturation dynamics compared to weak ones (Engelhard and  
229 Heino, 2004b; ICES, 2016). Whether cohort-to-cohort or year-to-year variability leads to prac-  
230 tically significant differences between estimations by cohort and by year is probably case-spe-  
231 cific.

232         Gulland's method also applies to irreversible life-history transitions other than matura-  
233 tion, such as metamorphosis, smolting, other ontogenetic niche shifts, and sex change. Each of  
234 these transitions might pose specific challenges, for example, in regard to the identification of

235 newly transitioned individuals. Furthermore, some of these transitions might not be strictly ir-  
236 reversible. For example, sex change can be bi-directional in some non-commercial reef fishes  
237 (e.g., Sunobe and Nakazono, 1993; Nakashima et al., 1996; Munday et al., 1998).

238 Skipped spawning has been implicated in a number of fish stocks (Rideout et al., 2005),  
239 including the stocks for which Gulland's method has been used (Engelhard and Heino, 2005;  
240 Skjæraasen et al., 2012). Skipped spawning has the potential to bias maturity ogives downwards,  
241 if not accounted for. For the documented levels of skipped spawning, this bias is modest, but  
242 may act in the same direction as other sources of bias. Furthermore, when good estimates of  
243 skipped spawning are available, the estimation of maturity ogives can readily be adjusted to  
244 account for it.

245 A major limitation of Gulland's method is that it requires data that are only seldom  
246 recorded: distinguishing first-time spawners from repeat spawners with routine observations is  
247 rarely possible. Exceptions mainly come from stocks with long spawning migrations (such as  
248 cod and herring) or an anadromous life cycle (such as shortnose sturgeon and steelhead trout).  
249 These show a so-called spawning check, a visually detectable change in the zonation pattern of  
250 otoliths or scales. Long migrations increase the energetic cost of spawning, making the slowing  
251 down of growth upon maturation more marked than it would otherwise be (e.g., Folkvord et al.,  
252 2014).

253 For Northeast Arctic cod, the recording of spawning checks is part of routine data col-  
254 lection (Mjanger et al., 2010). However, a strict validation of the assumption that the formation  
255 of the first spawning check is associated with the first spawning is as yet lacking. Experiments  
256 support the notion that substantial energy investments in reproduction are reflected by reduced  
257 otolith growth, but leave open the question whether such changes in the otolith growth of wild  
258 cod are always associated with reproduction or can originate also from other sources of ener-

259 genetic stress. Furthermore, it remains possible that some spawning cod do not develop a corre-  
260 sponding spawning check (Irgens, 2018; Arild Folkvord, University of Bergen, pers. comm.).  
261 Our numerical tests suggest that Gulland's method is not overly sensitive to moderate error  
262 rates in assessing whether a spawning fish is a first-time or second-time spawner.

263         There are a number of possibilities to identify first-time spawners in the absence of  
264 visually marked changes in otoliths or scales. First, numerical methods might allow detecting  
265 changes in growth that are not visually obvious as spawning checks (Rijnsdorp and Storbeck,  
266 1995; Engelhard et al., 2003; Baulier and Heino, 2008; Brunel et al., 2013). This requires back-  
267 calculations of growth, which can be based on both archived and fresh materials. The efficiency  
268 of such estimates could potentially be improved using automated image analyses. Unfortunately,  
269 detecting maturation from growth trajectories suffers from limited accuracy, especially for in-  
270 dividuals captured soon after maturation (Baulier and Heino, 2008; Brunel et al., 2013).

271         Second, advances in sclerochronology offer new possibilities for extracting life-history  
272 information from otoliths and scales that go far beyond traditional visual examinations. For  
273 example, patterns of ontogenetic vertical migrations in deep-sea fish can be deduced from oto-  
274 lith microstructures and stable-isotope composition (Lin et al., 2012; Chang et al., 2015). Like-  
275 wise, migrations of eel between marine, brackish, and freshwater environments can be deduced  
276 from the ratios of strontium and calcium in their otoliths (Jessop et al., 2008). An experimental  
277 study with European plaice (*Pleuronectes platessa*) showed that reproduction alters the zinc-  
278 to-calcium ratio in the blood plasma and otoliths of females, offering a potential tool for detect-  
279 ing both age at first reproduction and skipped spawning seasons (Sturrock et al., 2015). Also  
280 these methods apply to archived as well as fresh materials.

281         Third, histological methods offer what is potentially the most precise approach to sepa-  
282 rating first-time and repeat-spawning female fish, at least for determinately spawning boreal  
283 species: mature females with postovulatory follicles (POFs) are repeat-spawners, whereas those

284 lacking POFs are first-time spawners, provided that POFs persist long enough after spawning  
285 relative to the timing of the sample collection (Rideout et al., 2005). This condition is easily  
286 fulfilled in cod where POFs persist for more than a year (Witthames et al., 2010; Folkvord et  
287 al., 2014). Unfortunately, histological methods require samples that are specifically conserved  
288 for such analysis, which makes them unlikely to be applicable to historical materials. Moreover,  
289 obtaining age estimates still requires the reading of otoliths or scales.

290 Gulland's method offers the prospect of estimating maturity ogives for species in which  
291 it is difficult to obtain representative data on the juvenile part of the population. This will often  
292 necessitate collecting new kinds of data, or using existing materials in novel ways, in order to  
293 differentiate between first-time and repeat spawners. At the same time, elucidating individual  
294 life cycles with such new data will probably be valuable in its own right. The prospect of ob-  
295 taining information on a population's demographic composition by applying Gulland's method  
296 can add motivation for investing scarce resources in enhanced analyses of otoliths and scales,  
297 as well as into histology. Exciting options for future applications of the generalized Gulland's  
298 method presented here include obtaining information on the oceanic phase of anadromous it-  
299 eroparous salmonids and estimating the maturity ogives of enigmatic deep-sea fishes such as  
300 orange roughy.

### 301 **Acknowledgements**

302 We thank Arild Folkvord for discussions on cod otoliths and two reviewers for valuable feed-  
303 back that has helped us to improve the manuscript. MH acknowledges funding during the final  
304 preparation of this manuscript by the Ministry of Science and Technology (MOST), R.O.C.,  
305 grant number 105-2811-M-002-068.

306 **References**

- 307 Baulier, L., Heino, M., 2008. Norwegian spring-spawning herring as the test case of piece-  
308 wise linear regression method for detecting maturation from growth patterns. *J. Fish*  
309 *Biol.* 73, 2452–2467. <https://doi.org/10.1111/j.1095-8649.2008.02088.x>
- 310 Beverton, R.J.H., Holt, S.J., 1957. *On the Dynamics of Exploited Fish Populations*, Fishery  
311 Investigations Series. Ministry of Agriculture, Fisheries and Food, London.
- 312 Brunel, T., Ernande, B., Mollet, F.M., Rijnsdorp, A.D., 2013. Estimating age at maturation  
313 and energy-based life-history traits from individual growth trajectories with nonlinear  
314 mixed-effects models. *Oecologia* 172, 631–643. [https://doi.org/10.1007/s00442-012-](https://doi.org/10.1007/s00442-012-2527-1)  
315 [2527-1](https://doi.org/10.1007/s00442-012-2527-1)
- 316 Chang, N.-N., Liu, E.Y., Liao, Y.C., Shiao, J.-C., 2015. Vertical habitat shift of viviparous  
317 and oviparous deep-sea cusk eels revealed by otolith microstructure and stable-isotope  
318 composition. *J. Fish Biol.* 86, 845–853. <https://doi.org/10.1111/jfb.12605>
- 319 Devold, F., 1938. The North Atlantic halibut and net fishing. *Fisk. Skr. Ser. Havunders.* 5, 5–  
320 47.
- 321 Engelhard, G.H., Dieckmann, U., Godø, O.R., 2003. Age at maturation predicted from routine  
322 scale measurements in Norwegian spring-spawning herring (*Clupea harengus*) using  
323 discriminant and neural network analyses. *ICES J. Mar. Sci.* 60, 304–313.  
324 [https://doi.org/10.1016/S1054-3139\(03\)00017-1](https://doi.org/10.1016/S1054-3139(03)00017-1)
- 325 Engelhard, G.H., Heino, M., 2005. Scale analysis suggests frequent skipping of the second re-  
326 productive season in Atlantic herring. *Biol. Lett.* 1, 172–175.  
327 <https://doi.org/10.1098/rsbl.2004.0290>
- 328 Engelhard, G.H., Heino, M., 2004a. Maturity changes in Norwegian spring-spawning herring  
329 *Clupea harengus*: compensatory or evolutionary responses? *Mar. Ecol. Prog. Ser.* 272,  
330 245–256. <https://doi.org/10.3354/meps272245>

331 Engelhard, G.H., Heino, M., 2004b. Maturity changes in Norwegian spring-spawning herring  
332 before, during, and after a major population collapse. *Fish. Res.* 66, 299–310.  
333 [https://doi.org/10.1016/S0165-7836\(03\)00195-4](https://doi.org/10.1016/S0165-7836(03)00195-4)

334 Flores, A., Wiff, R., Díaz, E., 2015. Using the gonadosomatic index to estimate the maturity  
335 ogive: application to Chilean hake (*Merluccius gayi gayi*). *ICES J. Mar. Sci.* 72, 508–  
336 514. <https://doi.org/10.1093/icesjms/fsu155>

337 Folkvord, A., Jørgensen, C., Korsbrekke, K., Nash, R.D.M., Nilsen, T., Skjæraasen, J.E.,  
338 2014. Trade-offs between growth and reproduction in wild Atlantic cod. *Can. J. Fish.*  
339 *Aquat. Sci.* 71, 1106–1112. <https://doi.org/10.1139/cjfas-2013-0600>

340 Francis, R.I.C.C., Horn, P.L., 1997. Transition zone in otoliths of orange roughy (*Hoploste-*  
341 *thus atlanticus*) and its relationship to the onset of maturity. *Mar. Biol.* 129, 681–687.  
342 <https://doi.org/10.1007/s002270050211>

343 Gulland, J.A., 1969. *Manual of Methods for Fish Stock Assessment - Part 1. Fish Population*  
344 *Analysis* (No. M4), FAO Manuals in Fisheries Science. FAO, Rome.

345 Gulland, J.A., 1964. The abundance of fish stocks in the Barents Sea. *Rapp. Procès-Verbaux*  
346 *La Réunion. Cons. Perm. Int. Pour Explor. Mer* 155, 126–137.

347 Heino, M., Dieckmann, U., Godø, O.R., 2002. Estimating reaction norms for age and size at  
348 maturation with reconstructed immature size distributions: a new technique illustrated  
349 by application to Northeast Arctic cod. *ICES J. Mar. Sci.* 59, 562–575.  
350 <https://doi.org/10.1006/jmsc.2002.1192>

351 Hixon, M.A., Johnson, D.W., Sogard, S.M., 2014. BOFFFFs: on the importance of conserv-  
352 ing old-growth age structure in fishery populations. *ICES J. Mar. Sci.* 71, 2171–2185.  
353 <https://doi.org/10.1093/icesjms/fst200>

354 ICES, 2017. Report of the Arctic Fisheries Working Group (AFWG), 19–25 April 2017, Co-  
355 penhagen, Denmark (ICES Document No. CM 2017/ACOM:06). ICES, Copenhagen.



356 ICES, 2016. Report of the Benchmark Workshop on Pelagic stocks (WKPELA), 29 Febru-  
357 ary–4 March 2016, ICES Headquarters, Copenhagen, Denmark (ICES Document No.  
358 CM 2016/ACOM:34). ICES, Copenhagen.

359 ICES, 2008. Report of the Workshop on Maturity Ogive Estimation for Stock Assessment  
360 (WKMOG), 3–6 June 2008, Lisbon, Portugal (ICES Document No. CM  
361 2008/ACOM:33). ICES, Copenhagen.

362 Irgens, C., 2018. Otolith structure as indicator of key life history events in Atlantic cod (*Ga-*  
363 *dus morhua*) (PhD thesis). University of Bergen, Bergen, Norway.

364 Jessop, B.M., Cairns, D.K., Thibault, I., Tzeng, W.N., 2008. Life history of American eel *An-*  
365 *guilla rostrata*: new insights from otolith microchemistry. *Aquat. Biol.* 1, 205–216.  
366 <https://doi.org/10.3354/ab00018>

367 Jørgensen, C., Ernande, B., Fiksen, Ø., Dieckmann, U., 2006. The logic of skipped spawning  
368 in fish. *Can. J. Fish. Aquat. Sci.* 63, 200–211.

369 Jørgensen, T., 1990. Long-term changes in age at sexual maturity of Northeast Arctic cod  
370 (*Gadus morhua* L.). *J. Cons. Int. Pour Explor. Mer* 46, 235–248.  
371 <https://doi.org/10.1093/icesjms/46.3.235>

372 Köster, F.W., Trippel, E.A., Tomkiewicz, J., 2013. Linking size and age at sexual maturation  
373 to body growth, productivity and recruitment of Atlantic cod stocks spanning the  
374 North Atlantic. *Fish. Res.* 138, 52–61. <https://doi.org/10.1016/j.fishres.2012.07.002>

375 Lea, E., 1928. Undersøkelser over den norske sild. Fra umoden til moden. Årsberet.  
376 Vedkomm. Nor. Fisk. 4, 1–36.

377 Lin, H.-Y., Shiao, J.-C., Chen, Y.-G., Iizuka, Y., 2012. Ontogenetic vertical migration of  
378 grenadiers revealed by otolith microstructures and stable isotopic composition. *Deep-*  
379 *Sea Res. Part Oceanogr. Res. Pap.* 61, 123–130.  
380 <https://doi.org/10.1016/j.dsr.2011.12.005>

381 Mjanger, H., Hestenes, K., Svendsen, B.V., Wenneck, T. de L., 2010. Håndbok for  
382 prøvetaking av fisk og krepsdyr. Versjon 3.16. August 2010. Institute of Marine  
383 Research, Bergen, Norway.

384 Munday, P.L., Caley, M.J., Jones, G.P., 1998. Bi-directional sex change in a coral-dwelling  
385 goby. *Behav. Ecol. Sociobiol.* 43, 371–377. <https://doi.org/10.1007/s002650050504>

386 Murua, H., Saborido-Rey, F., 2003. Female reproductive strategies of marine fish species of  
387 the North Atlantic. *J. Northwest Atl. Fish. Sci.* 33, 23–31.

388 Nakashima, Y., Kuwamura, T., Yogo, Y., 1996. Both-ways sex change in monogamous coral  
389 gobies, *Gobiodon* spp. *Environ. Biol. Fishes* 46, 281–288.  
390 <https://doi.org/10.1007/BF00005004>

391 Narver, D.W., 1969. Age and size of steelhead trout in the Babine River, British Columbia. *J.*  
392 *Fish. Res. Board Can.* 26, 2754–2760. <https://doi.org/10.1139/f69-269>

393 Ottersen, G., Hjermann, D.Ø., Stenseth, N.C., 2006. Changes in spawning stock structure  
394 strengthen the link between climate and recruitment in a heavily fished cod (*Gadus*  
395 *morhua*) stock. *Fish. Oceanogr.* 15, 230–243.

396 Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish popula-  
397 tions, Bulletin of the Fisheries Research Board of Canada. Department of the Environ-  
398 ment, Fisheries and Marine Service, Ottawa, Canada.

399 Rideout, R.M., Rose, G.A., Burton, M.P.M., 2005. Skipped spawning in female iteroparous  
400 fishes. *Fish Fish.* 6, 50–72.

401 Rijnsdorp, A.D., Storbeck, F., 1995. Determining the onset of sexual maturity from otoliths of  
402 individual North Sea plaice, *Pleuronectes platessa* L., in: Secor, D.H., Dean, J.M.,  
403 Campana, S.E. (Eds.), *Recent Developments in Fish Otolith Research*, The Belle W.  
404 Baruch Library in Marine Science. University of South Carolina Press, pp. 581–598.

405 Rollefson, G., 1933. The otoliths of the cod. *Fisk. Skr. Ser. Havunders.* 4, 3–14.

- 406 Runnström, S., 1936. A study on the life history and migrations of the Norwegian spring-her-  
407 ring based on the analysis of the winter rings and summer zones of the scale. Fisk.  
408 Skr. Ser. Havunders. 5, 1–107.
- 409 Skjæraasen, J.E., Nash, R.D.M., Korsbrekke, K., Fonn, M., Nilsen, T., Kennedy, J., Nedreaas,  
410 K.H., Thorsen, A., Witthames, P.R., Geffen, A.J., Høie, H., Kjesbu, O.S., 2012. Fre-  
411 quent skipped spawning in the world's largest cod population. Proc. Natl. Acad. Sci.  
412 U. S. A. 109, 8995–8999. <https://doi.org/10.1073/pnas.1200223109>
- 413 Stearns, S.C., Crandall, R.E., 1984. Plasticity for age and size at sexual maturity: a life-history  
414 response to unavoidable stress, in: Potts, G.W., Wootton, R.J. (Eds.), Fish Reproduc-  
415 tion: Strategies and Tactics. Academic Press, London, pp. 13–33.
- 416 Sturrock, A.M., Hunter, E., Milton, J.A., Johnson, R.C., Waring, C.P., Trueman, C.N., 2015.  
417 Quantifying physiological influences on otolith microchemistry. Methods Ecol. Evol.  
418 6, 806–816. <https://doi.org/10.1111/2041-210X.12381>
- 419 Sunobe, T., Nakazono, A., 1993. Sex change in both directions by alteration of social domi-  
420 nance in *Trimma okinawae* (Pisces: Gobiidae). Ethology 94, 339–345.  
421 <https://doi.org/10.1111/j.1439-0310.1993.tb00450.x>
- 422 Svåsand, T., Ajiad, A.M., Buehler, V., Carvalho, G.R., Clemmesen, C., Dahle, G., Hard, J.J.,  
423 Hauser, L., Hutchinson, W.F., Jakobsen, T., Kjesbu, O.S., Moksness, E., Otterå, H.,  
424 Paulsen, H., Schnack, D., Solemdal, P., Thorsen, A., 2003. Demonstration of maternal  
425 effects of Atlantic cod: combining the use of unique mesocosm and novel molecular  
426 techniques (MACOM). Final Report, Project QLK5-CT1999-01617. Institute of Ma-  
427 rine Research, Bergen, Norway.
- 428 Taubert, B.D., 1980. Reproduction of shortnose sturgeon (*Acipenser brevirostrum*) in Ho-  
429 lyoke Pool, Connecticut River, Massachusetts. Copeia 1980, 114–117.
- 430 Trippel, E.A., 1995. Age at maturity as stress indicator in fisheries. BioScience 45, 759–771.

431 <https://doi.org/10.2307/1312628>

432 Trippel, E.A., Kjesbu, O.S., Solemdal, P., 1997. Effects of adult age and size structure on re-  
433 productive output in marine fishes, in: Chambers, R.C., Trippel, E.A. (Eds.), Early  
434 Life History and Recruitment in Fish Populations. Chapman and Hall, London, pp.  
435 63–102.

436 Witthames, P.R., Thorsen, A., Kjesbu, O.S., 2010. The fate of vitellogenic follicles in experi-  
437 mentally monitored Atlantic cod *Gadus morhua* (L.): Application to stock assessment.  
438 Fish. Res. 104, 27–37. <https://doi.org/10.1016/j.fishres.2009.11.008>

439 Yaragina, N.A., 2010. Biological parameters of immature, ripening, and non-reproductive,  
440 mature northeast Arctic cod in 1984–2006. ICES J. Mar. Sci. 67, 2033–2041.  
441 <https://doi.org/10.1093/icesjms/fsq059>

442 Zuykova, N.V., Koloskova, V.P., Mjanger, H., Nedreaas, K.H., Senneset, H., Yaragina, N.A.,  
443 Aagotnes, P., Aanes, S., 2009. Age determination of Northeast Arctic cod otoliths  
444 through 50 years of history. Mar. Biol. Res. 5, 66–74.  
445 <https://doi.org/10.1080/17451000802454874>

446

447 **Figure captions**

448 **Fig. 1.** Schematic illustration of Gulland's method when survival is equal for all individuals at  
449 a given age. At any point in time, a cohort is composed of juvenile (J), first-spawning (F), and  
450 repeat-spawning (R) individuals. For all ages  $t$ , the proportion of repeat spawners among all  
451 spawners is known from sampling,  $r_t = n_t^R / (n_t^F + n_t^R)$ . The maturity ogive  $o_t$  is known from  
452 sampling to equal 1 down to the latest age  $T$  at which first-time spawners are still present in the  
453 samples. For earlier ages  $t < T$ ,  $o_t$  can be calculated iteratively using the equation  $o_{t-1} = o_t r_t$ .  
454 For understanding Gulland's method, it is important to recognize that the proportions  $o_t$  of ma-  
455 ture individuals among all individuals and the proportions  $r_t$  of repeat spawners among all ma-  
456 ture individuals are measured relative to different totals (all individuals vs. all mature individ-  
457 uals), as indicated by the gray extensions of the black curly braces.

458 **Fig. 2.** Sensitivity of the estimated proportion of mature individuals to variations in relative  
459 survival between maturity stages. The left and right columns correspond to situations that could  
460 be encountered when estimating maturity proportions for near median and relatively early ages,  
461 respectively, that is, relatively high and low proportions of mature and first-spawning individ-  
462 uals. On the left, assuming that all survival ratios are equal, the estimated maturity proportion  
463 is  $o_{t-1} = o_t r_t \approx 0.56$ , whereas on the right, it is much lower, at  $o_{t-1} \approx 0.062$ . The color bands  
464 indicate how much this estimate deviates from the true value (red: overestimation; blue: under-  
465 estimation). Each color band has a width of 0.05. The white band is centered on the true value;  
466 hence, it covers an area in which the absolute error is smaller than 2.5%. In the top row, all  
467 survival ratios are varied, but  $s_{t-1}^{JJ} = s_{t-1}^{JF}$  are kept equal, whereas in the middle row,  $s_{t-1}^{JF} =$   
468  $s_{t-1}^{FR}$  are kept equal. In the bottom row,  $s_{t-1}^{JJ}$  and  $s_{t-1}^{JF}$  are varied independently, while  $s_{t-1}^{FR} = 1$   
469 is kept fixed. Notice that all axes are logarithmic.

470 **Fig. 3.** Application of the generalized Gulland's method to the 1928 cohort of Northeast Arctic

471 cod, *Gadus morhua*. In (a), maturing fish and first-time spawners are assumed to have experi-  
 472 enced the same survival probability as repeat spawners ( $s_t^{JF}/s_t^{RR} = s_t^{FR}/s_t^{RR} = 1$ ), whereas in  
 473 (b), they are assumed to suffer from reduced survival during the spawning migration  
 474 ( $s_t^{JF}/s_t^{RR} = 0.7$  and  $s_t^{FR}/s_t^{RR} = 0.9$ ). In both panels, the ratio of juvenile to repeat spawner sur-  
 475 vival probability is varied; the size of the circular markers increases with the survival ratios  
 476  $s_t^{JJ}/s_t^{RR}$ , which apply to all ages  $t$ . Filled circles correspond to the simpler case in which sur-  
 477 vival is independent of maturation stage, as is assumed when using the original Gulland's  
 478 method without our generalization. The results show how estimation errors as large as 100%  
 479 can result when applying the original Gulland's method to situations in which age-specific sur-  
 480 vival probabilities are affected by maturation stage.

481 **Fig. 4.** Application of the generalized Gulland's method to the 1928 cohort of Northeast Arctic  
 482 cod when accounting for (a) skipped spawning and (b) errors in assessing an individual's  
 483 spawning experience. When some mature fish skip the spawning migration, samples from the  
 484 spawning grounds show too low proportions  $r_T$  of repeat spawners among all mature individu-  
 485 als. In (a),  $r_T$  is corrected for this underrepresentation as  $r'_T = r_T(1 + x)/[r_T(1 + x) + 1 - r_T]$ ,  
 486 where  $x$  is the proportion of mature fish skipping spawning. In (b), it is assumed that there is a  
 487 20% probability of assigning a first-time spawner as a second-time spawner, and vice versa.  
 488 The thick curve shows the true ogive when spawning experience is assumed to have been cor-  
 489 rectly estimated. The boxes show the median value together with the interquartile range of 1000  
 490 Monte-Carlo replicates in which erroneous assignments are present. Whiskers extend to the  
 491 most extreme data point no more than 1.5 times the interquartile range away from the box, and  
 492 dots show more extreme data points.

Fig. 1





Fig. 2  $o_t = 0.75, r_t = 0.75$



$o_t = 0.25, r_t = 0.25$











Fig. 3  $s^{JJ}/s^{RR} = 0.5...2$ ,  $s^{JF}/s^{RR} = 1$ ,  $s^{FR}/s^{RR} = 1$



$s^{JJ}/s^{RR} = 0.5...2$ ,  $s^{JF}/s^{RR} = 0.7$ ,  $s^{FR}/s^{RR} = 0.9$





Fig. 4

**Skipped spawning****Errors in spawning experience**