



Supplement of

Quantifying memory and persistence in the atmosphere–land and ocean carbon system

Matthias Jonas et al.

Correspondence to: Matthias Jonas (jonas@iiasa.ac.at)

The copyright of individual parts of the supplement might differ from the article licence.

Supplementary Information (SI)

SI1: To equation (2)

With $\dot{\varepsilon}(t) = \alpha \exp(\alpha t)$ and $\alpha > 0$, $\sigma(0) = 0$ and $\beta = 1 + \alpha \frac{D}{K}$:

$$\begin{aligned}\sigma(t) &= K \int_0^t \alpha \exp(\alpha \tau) \exp\left(\frac{K}{D}(\tau - t)\right) d\tau = K \alpha \exp\left(-\frac{K}{D}t\right) \int_0^t \exp\left(\frac{K}{D}\beta \tau\right) d\tau \\ &= \frac{D}{\beta} \alpha \exp\left(-\frac{K}{D}t\right) \left(\exp\left(\frac{K}{D}\beta t\right) - 1\right) = \frac{D}{\beta} \alpha \exp(\alpha t) \left(1 - \exp\left(-\frac{K}{D}\beta t\right)\right) \\ &= \frac{D}{\beta} \dot{\varepsilon}(t) (1 - q_\beta^t)\end{aligned}$$

where $q_\beta^t = \exp\left(-\frac{K}{D}\beta t\right)$. Introducing the dimensionless time $n = \frac{t}{\Delta t}$ (where $\Delta t = 1 \text{ yr}$), equation (2) takes the form

$$\sigma(n) = \frac{D}{\beta} \frac{\alpha_n}{\Delta t} \exp(\alpha_n n) (1 - q_\beta^n) = \frac{D}{\beta} \dot{\varepsilon}(n) (1 - q_\beta^n) \quad (\text{SI1-1})$$

where $\alpha_n := \alpha \Delta t$, $q_\beta = q_\alpha q$, $q_\alpha = \exp(-\alpha \Delta t)$, $q = \exp\left(-\frac{K}{D}\Delta t\right)$, and

$$q_\beta^t = \left(\exp\left(-\frac{K}{D}\beta \Delta t\right)\right)^n.$$

SI2: To equation (3)

We start from equation (2b) in the form $\sigma_D(q, n) := \frac{1}{D} \sigma(n) = \dot{\varepsilon}(n) \frac{\ln q}{\ln q - \alpha_n} (1 - q_\alpha^n q^n)$ with

$$\frac{1}{\beta} = \frac{\ln q}{\ln q - \alpha_n}:$$

$$\begin{aligned}q \frac{\partial}{\partial q} \sigma_D(q, n) &= \dot{\varepsilon}(n) q \left\{ \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} \right) (1 - q_\beta^n) - \frac{\ln q}{\ln q - \alpha_n} q_\alpha^n \frac{\partial}{\partial q} q^n \right\} \\ &= \dot{\varepsilon}(n) \left\{ q \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} \right) (1 - q_\beta^n) - \frac{1}{\beta} q_\beta^n n \right\}\end{aligned} \quad (\text{SI2-1a})$$

where we can avoid the effort of writing out the 1st derivative on the right side; and

$$q q_\alpha^n \frac{\partial}{\partial q} q^n = q q_\alpha^n q^{n-1} n = q_\beta^n n.$$

On the other hand, with $S_n = \frac{1-q_\beta^n}{1-q_\beta}$ and the help of equation (SI3-4a):

$$\begin{aligned}
q \frac{\partial}{\partial q} \sigma_D(q, n) &= \dot{\varepsilon}(n) q \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} (1 - q_\beta) S_n \right) \\
&= \dot{\varepsilon}(n) q \left\{ \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} \right) (1 - q_\beta) S_n - \frac{1}{\beta} \frac{\partial q_\beta}{\partial q} S_n + \frac{1}{\beta} (1 - q_\beta) \frac{\partial q_\beta}{\partial q} \frac{\partial S_n}{\partial q_\beta} \right\} \\
&= \dot{\varepsilon}(n) \left\{ q \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} \right) (1 - q_\beta) S_n - \frac{1}{\beta} q_\beta S_n + \frac{1}{\beta} (1 - q_\beta) S_n \frac{q_\beta}{S_n} \frac{\partial S_n}{\partial q_\beta} \right\} \\
&= \dot{\varepsilon}(n) \left\{ q \frac{\partial}{\partial q} \left(\frac{\ln q}{\ln q - \alpha_n} \right) (1 - q_\beta^n) - \frac{1}{\beta} q_\beta S_n + \frac{1}{\beta} (1 - q_\beta^n) T \right\}
\end{aligned} \tag{SI2-1b}$$

Balancing equations (SI2-1a) and (SI2-1b) yields equation (3):

$$T = -\frac{q_\beta^n}{1 - q_\beta^n} n + \frac{q_\beta}{1 - q_\beta}. \tag{SI2-2}$$

$T = T(q, n)$ is a characteristic function of the Maxwell body (MB).

SI3: Justifying T as delay time

We can let any function f of time t (dimensionless throughout SI3 and $\in \mathbb{N}_0$ without restricting generality) depend increasingly on previous times by applying the approach of a simple weighted average and a weighting fading away exponentially backward in time ($q < 1$):

$$\begin{aligned}
y_1(t) &= f\left(\frac{q^0 t}{q^0}\right) \\
y_2(t) &= f\left(\frac{q^0 t + q^1 (t-1)}{q^0 + q^1}\right) \\
&\dots \\
y_k(t) &= f\left(\frac{q^0 t + q^1 (t-1) + q^2 (t-2) + \dots + q^{k-1} (t-(k-1))}{\sum_{i=0}^{k-1} q^i}\right) = f(t - T)
\end{aligned} \tag{SI3-1}$$

($t \geq k \in \mathbb{N}_0$) with T appearing as delay time in the argument of the function f . The denominator of the argument in the middle is given by

$$S_k = \sum_{i=0}^{k-1} q^i = \frac{1-q^k}{1-q}; \quad (\text{SI3-2})$$

while the numerator can be transformed with the help of¹

$$\sum_{k=a}^{b-1} k^m z^k = \left(z \frac{d}{dz} \right)^m \frac{z^b - z^a}{z-1} \quad (z \neq 1),$$

here with i instead of k , and q instead of z , and $a = 0$, $b = k$, and $m = 1$

$$\sum_{i=0}^{k-1} i q^i = q \frac{d}{dq} \frac{q^k - q^0}{q-1} = q \frac{d}{dq} \frac{1-q^k}{1-q} = q \frac{d}{dq} S_k \quad (q \neq 1) \quad (\text{SI3-3})$$

to derive T :

$$T = \frac{q}{S_k} \frac{d}{dq} S_k. \quad (\text{SI3-4a})$$

Similar to and in accordance with equation (3), carrying out the derivation by q on the right side yields

$$T = \frac{q}{S_k} \frac{1}{1-q} \left(-q^{k-1} k + S_k \right) = -\frac{q^k}{1-q^k} k + \frac{q}{1-q}. \quad (\text{SI3-4b})$$

It is straightforward to show by applying l'Hospital that $\lim_{k \rightarrow \infty} (q^k k) = 0$. Thus:

$$T_{k \rightarrow \infty} (= T_\infty) = \frac{q}{1-q} = q \frac{1}{1-q} = q S_{k \rightarrow \infty}. \quad (\text{SI3-5})$$

To strengthen the justification of T as delay time for the exponential function $y(t) = 1 - \exp(-ct) = 1 - q^t$ with $\ln(q) = -c$ and $c > 0$, it is useful to consider the power-

law case $y(t) = ct^q$. Here, the ratio $\frac{y}{\dot{y}}$ with $T = q$ functions as a linearizer such that

$$\frac{y - \dot{y} b}{\dot{y}} = a(t - T) \Leftrightarrow \frac{y - \dot{y}}{\dot{y}} = \frac{1}{T}(t - T); \text{ where } b = aT \text{ is the intercept, } T = \frac{\dot{y}}{y} t \text{ is the}$$

intersection with the time axis, and the difference $t - T$ can be expressed as well as weighted

$$(w) \text{ (or moving weighted) average } (t - T) = \frac{\sum_{i=0}^{k-1} w_{k-i} (t - i)}{\sum_{i=0}^{k-1} w_{k-i}}. \text{ } T \text{ being constant is in}$$

line with the finding (not shown here) that the change in memory can be considered Gaussian backward in time.

Similar for the exponential function $y(t) = 1 - q^t$. Here, q and t appear mirrored to the power-law case. Nonetheless, T (reduced by T_∞) in equation (SI3-5) can also be expressed, in principle (i.e., apart from additional factors), by the operation

$$T_{\text{red}} = T - T_{\infty} = \frac{1}{\ln(q)} \frac{\dot{y}}{y} t. \quad (\text{SI3-6})$$

However, despite this agreement, the change in memory here is exponential backward in time.

Equation (SI3-6) generalizes to $T - T_{\infty} = \frac{1}{\ln(q_{\beta})} \frac{\dot{\sigma} - \alpha \sigma}{\sigma} t$ in the case of equation (2a).

SI4: To equation (4) reflecting the history of the MB

Rewriting equation (4) shows that it reflects the history of the MB:

$$\begin{aligned} S_n &= \frac{q_{\beta}^n - 1}{q_{\beta} - 1} = \frac{1}{q_{\beta} - 1} \left\{ (q_{\beta}^n - q_{\beta}^{n-1}) + (q_{\beta}^{n-1} - q_{\beta}^{n-2}) + q_{\beta}^{n-2} + \dots + q_{\beta} + (q_{\beta} - 1) \right\} \\ &= \frac{1}{q_{\beta} - 1} \left\{ q_{\beta}^{n-1} (q_{\beta} - 1) + q_{\beta}^{n-2} (q_{\beta} - 1) + \dots + q_{\beta}^0 (q_{\beta} - 1) \right\} = \sum_{i=0}^{n-1} q_{\beta}^i = \text{Past} \end{aligned} \quad (\text{SI4-1})$$

SI5: To monitoring $\ln(M \cdot P)$

According to equations (3)–(5):

$$M_{\infty} = \frac{1}{1 - q_{\beta}} = \frac{T_{\infty}}{q_{\beta}} \quad \text{and} \quad P_{\infty} = \frac{1}{T_{\infty}}. \quad (\text{SI5-1,2})$$

Hence:

$$\frac{1}{M_{\infty} P_{\infty}} = q_{\beta} = \exp\left(-\left(\frac{K}{D} + \alpha\right)\Delta t\right) = \exp\left(-\frac{K}{D}\beta\Delta t\right) \Leftrightarrow \ln(M_{\infty} P_{\infty}) = \frac{K}{D}\beta\Delta t = \lambda_{\beta} = \lambda\beta$$

with q_{β} and q as defined under Methods, and $\lambda_{\beta} = \lambda\beta$ with $\lambda = \frac{K}{D}\Delta t$. Thus, the ratio $\frac{\lambda}{\ln(MP)}$

allows indicating how much smaller the system's natural rate of change in the numerator turns out compared to the system's rate of change in the denominator under continued increase in stress. This gradual build-up relative to λ (with K/D constant) is limited by β^{-1} .

SI6: Overview of data and conversion factors

Tab. SI6-1: Overview of the data used in the paper. All data refer to the global scale (or are assumed to be globally representative).

Data	Data Source	Time range	Brief description
Atmospheric CO ₂ concentration (in ppm)	2 Degrees Institute, Canada ²	1750–1955	Ice core data (75-year smoothed); Law Dome, Antarctica
		1959–1979	Atmospheric measurements (annual means); Mouna Loa, Hawaii
	Global Monitoring Laboratory, NOAA, USA ³	1980–2018	
CO ₂ emissions from fossil-fuel combustion and cement production (in PgC yr ⁻¹)	Global Carbon Project ⁴	1751–1958	Global estimates derived from energy statistics by nation and year
		1959–2015	
Land-use change emissions (in PgC yr ⁻¹)		1850–1958	Global mean values derived from multiple models
		1959–2015	
Net primary production (in PgC yr ⁻¹)	O'Sullivan et al. (2019) ⁵	1900–2016	Model-based global mean values (Community Land Model; CLIM4.5-BGC)
Dissolved organic carbon (in $\mu\text{mol kg}^{-1}$)	Bates et. al. (2014) ⁶	1983–2012 (max. range)	Shipboard observations (annual means); from 7 sites (2 in the subpolar North Atlantic and 5 in the tropical/subtropical/temperate waters of the North Atlantic and Pacific)

Tab. SI6-2: Overview of the conversion factors used in the paper.

From	To	Value	Unit	Source
C	CO ₂	3.664	gCO ₂ (gC) ⁻¹	CDIAC (2012: Tab. 3) ⁷
ppmv CO ₂	PgC	2.120	PgC ppmv ⁻¹	Ciais et al. (2014: Tab. 6.1) ⁸
ppmv CO ₂	Pa	0.101325	Pa (10 ⁶ ppmv) ⁻¹	CDIAC (2012: Tab. 3) ⁷ and Dalton's law ⁹

SI7: Use of equation (9) to estimate the photosynthetic carbon flux ratio $\Delta\text{Ph}_i/\text{Ph}$

The leaf-level factor L denotes the relative leaf photosynthetic response to a 1 ppmv change in the atmospheric concentration of CO₂. The photosynthetic limits L_1 (photosynthesis limited by electron transport) and L_2 (photosynthesis limited by rubisco activity) are determined by using equations (7) and (9) in Luo et al. (1996).¹⁰

We follow equation (9) to derive the photosynthetic carbon flux ratio $\Delta\text{Ph}_i/\text{Ph}$ by the change in L_i , which we describe by means of a geometric sequence (with the common ratio $1-q_{L_i}$).

We demonstrate the quality of this approximation by comparing our results (to the extent possible) with those cited by Luo et al. (1996). Dropping index i :

$$\begin{aligned}
& L_{\text{high}} - L_{\text{low}} \\
& = \Delta L = L_{\text{high}} + L_{\text{high}}(1 - q_L) + \dots + L_{\text{high}}(1 - q_L)^{(\Delta\text{CO}_2 - 1)} \quad (\text{SI7-1}) \\
& = L_{\text{high}} \sum_{k=0}^{\Delta\text{CO}_2 - 1} (1 - q_L)^k = L_{\text{high}} \frac{1 - (1 - q_L)^{\Delta\text{CO}_2}}{1 - (1 - q_L)}
\end{aligned}$$

where $q_L = \Delta L / (L_{\text{high}} \Delta\text{CO}_2)$. (We follow the authors and express L_i in units of % [and not in % ppmv⁻¹]. To express q_L in units of 1, we consider ΔCO_2 dimensionless [equivalent to multiplying ΔCO_2 with ppmv⁻¹].) The term L_{high} has to be replaced by the term $L_{\text{high}} f_{\text{ppm}}$ if ΔL is not calculated per 1-ppmv step but per 1-year step (when the change in ppmv is not necessarily 1 ppmv; see also SD1). With the values in Table SI7-1, equation (SI7-1) allows accumulated ΔL_i values to be derived which can be compared with the $\Delta\text{Ph}_i/\text{Ph}$ values reported by Luo et al. (1996) in their Table 1.¹⁰ The agreement is sufficient for our purposes (Tab. SI7-2).

Tab. SI7-1: Limits of the relative leaf photosynthetic response to a 1 ppm change in the atmospheric concentration of CO₂ using equations (7) and (9) in Luo et al. (1996).

Time	CO ₂	L ₁	L ₂
yr	ppmv	%	%
preindustrial	280	0.1827	0.3520
1958	315	0.1457	0.2969
1992	355.5	0.1155	0.2495
1993	357	0.1146	0.2479

Tab. SI7-2: Comparison of ΔL_i (accumulated) derived with equation (SI7-1) with $\Delta\text{Ph}_i/\text{Ph}$ as listed in Table 1 in Luo et al. (1996).

Period	ΔCO_2	q_{L1}	q_{L2}	ΔL_1	ΔL_2	$\Delta\text{Ph}_i/\text{Ph}$
yr	ppmv	1	1	%	%	%
1992–1993	1.5	0.005358	0.004031	0.17	0.37	0.17–0.37
1958–1993	42	0.005080	0.003929	5.6	11.5	5.6–12.1
preindustrial –1993	77	0.004839	0.003840	11.8	23.5	11.8–25.5

SI8: The compression module referring to a tropospheric expansion of 20 m (standard atmosphere)

The standard atmosphere assigns a temperature gradient of -6.5 °C/1000 m up to the tropopause at 11 km. The isentropic coefficient of expansion γ varies with temperature and atmospheric CO₂ concentration: γ increases with decreasing T and decreases with increasing atmospheric CO₂.¹¹ However, in the case of dry air and no change in its chemical composition, the compression module K_{ad} can be expected to stay constant. Here we provide an overview of

the altitudes different isentropic coefficients of expansion refer to assuming a tropospheric expansion of 20 m;^{12,13} and, thereupon, determine K_{ad} .

Combining equations (19) and (20b):

$$K_{ad} = \gamma p = -\frac{\Delta p}{\Delta V/V} \quad (\text{SI8-1})$$

where the difference in pressure for a difference in altitude $\Delta h = h_2 - h_1$ is given by

$$\Delta p = p_2 - p_1 = p_0 \left[(1 - a(h_1 + \Delta h))^b - (1 - a h_1)^b \right]$$

according to equation (7) in Cavcar (2000)¹⁴ with $p_0 = 1013.25 \text{ hPa}$, $a = 0.0065/T_0$, $T_0 = 288.15 \text{ K}$, $b = 5.2561$, and h the altitude in units of meter;

and the difference in volume by

$$\frac{\Delta V}{V} = \frac{V_2 - V_1}{V_1 - V_{\text{Earth}}} = \frac{(r_{\text{Earth}} + (h_1 + \Delta h)/1000)^3 - (r_{\text{Earth}} + h_1/1000)^3}{(r_{\text{Earth}} + h_1/1000)^3 - r_{\text{Earth}}^3}$$

with $r_{\text{Earth}} = 6371 \text{ km}$.

Letting p refer to p_1 in equation (SI8-1) and solving for γ :

$$\gamma = \frac{1 - \left\{ (1 - a(h_1 + \Delta h)) / (1 - a h_1) \right\}^b}{\Delta V/V} \quad (\text{SI8-2})$$

Setting $\Delta h = 20 \text{ m}$ in agreement with observations, equation (SI8-2) allows calculating γ in dependence of h_1 (see Tab. SI8-1). As can also be seen from the table, the value of K_{ad} ranges between 400 and 412 hPa.

Tab. SI8-1: Standard atmosphere: isentropic coefficient of expansion γ and compression module K_{ad} for a tropospheric expansion of 20 m at different altitudes.

h_1	γ	p_1	K_{ad}
m	1	hPa	hPa
Input	Eq. (SI8-2)	Eq. (7) in Cavcar (2000)	Eq. (SI8-1)
7,100	1.000	404.8	404.8
7,685	1.100	372.5	409.6
8,255	1.200	343.0	411.6
8,810	1.300	316.2	411.1
8,865	1.310	313.6	411.0
9,345	1.400	292	408.8
9,360	1.403	291.3	408.7
9,865	1.500	269.9	404.9
10,370	1.600	249.7	399.6

SI9: Equation (1a) with strain given by a second-order polynomial

We start from $\varepsilon(t) = c_2 t^2 + c_1 t$. Inserting $\dot{\varepsilon}(t) = 2c_2 t$ into equation (1a) with $\sigma(0) = 0$ and

$$\int x e^{cx} dx = e^{cx} \frac{cx - 1}{c^2} :^{15}$$

$$\begin{aligned} \sigma(t) &= K \int_0^t \dot{\varepsilon}(\tau) \exp\left(\frac{K}{D}(\tau - t)\right) d\tau = 2c_2 K \exp\left(-\frac{K}{D}t\right) \int_0^t \tau \exp\left(\frac{K}{D}\tau\right) d\tau \\ &= 2c_2 K \exp\left(-\frac{K}{D}t\right) \left\{ \exp\left(\frac{K}{D}\tau\right) \frac{\frac{K}{D}\tau - 1}{\left(\frac{K}{D}\right)^2} \right\}_0^t = 2c_2 K \exp\left(-\frac{K}{D}t\right) \left\{ \exp\left(\frac{K}{D}t\right) \frac{\frac{K}{D}t - 1}{\left(\frac{K}{D}\right)^2} + \frac{1}{\left(\frac{K}{D}\right)^2} \right\} \\ &= 2c_2 \frac{D^2}{K} \left\{ \frac{K}{D}t - 1 + \exp\left(-\frac{K}{D}t\right) \right\} \xrightarrow{t \rightarrow \infty} 2c_2 D \left(t - \frac{D}{K}\right) \end{aligned}$$

SI10: Overview of parameters in experiments B and C

Table SI10-1 provides an overview of the parameters which result from the set of stress and strain explicit experiments B and C. They can be understood as a repetition of the 1959–2015 Case 0 experiment (see A.1 in the Results section), but with the difference that now upstream emissions as of 1900 (B) or 1850 (C), respectively, are considered; thus allowing initial conditions for 1959 other than zero as in the Case 0 experiment to be taken into account:

Case 0: 1959–2015

B: 1900–1958 (upstream emissions), 1959–2015

C: 1850–1958 (upstream emissions), 1959–2015.

The experiments are ordered consecutively in term of time. By way of contrast, Table SI10-2 comprises the parameters of the three 1959–2015 periods in the form of min–max intervals. Except for the exponential growth factor α , these intervals are dominated by Case 0 and B (1959–2015) parameters (as shown by the background color of the cells); mirroring the fact that we had difficulties with describing the entire upstream period 1850–1958 by means of a single exponential growth factor (0.0151 yr⁻¹).

Nonetheless, Table SI10-2 allows drawing a number of robust results:

- The compression modulus K increased between 1850 and 1959–2015 from ~2 to 10–13 Pa (the atmosphere became less compressible);
- while the damping constant D decreased between 1850 and 1959–2015 from ~468 to 459–462 Pa yr (the uptake of carbon by land and oceans became less viscous);
- with the consequence that the ratio $\lambda = K/D$ increased between 1850 and 1959–2015 from ~0.004–0.005 yr⁻¹ to 0.021–0.028 yr⁻¹ (i.e., by a factor of 4 to 6).
- Delay time T_∞ decreased (hence persistence P_∞ increased) between 1850 and 1959–2015 from ~51 (~0.02) to 18–21 (0.047–0.055) on the dimensionless timescale;

- while memory M_∞ decreased between 1850 and 1959–2015 from ~52 to 19–22 on the dimensionless timescale.

Tab. SI10-1: Overview of parameters in experiments B and C.

Parameters		Case 0	B		C	
		1959–2015	1900–1958	1959–2015	1850–1958	1959–2015
stress explicit						
$\sigma(0)$	Pa	0	0	5.8	0	7.8
K	Pa	9.9	2.4	12.7	2.1	11.6
D	Pa yr	461.5	467.7	459.2	467.9	460.1
$\lambda^{a,b}$	yr ⁻¹	0.0214	0.0051	0.0276	0.0045	0.0253
λ^{-1}	yr	46.8	196.3	36.3	223.5	39.6
α^a	yr ⁻¹	0.0247	0.0228	0.0262	0.0151	0.0281
β	1	2.158	5.475	1.951	4.371	2.112
λ_β^a	yr ⁻¹	0.0461	0.0279	0.0538	0.0196	0.0533
λ_β^{-1}	yr	21.7	35.9	18.6	51.1	18.7
q_β	1	0.9549	0.9725	0.9476	0.9806	0.9481
T_∞	1	21.2	35.4	18.1	50.6	18.3
M_∞ = T_∞/q_β	1	22.2	36.4	19.1	51.6	19.3
P_∞ = $1/T_\infty$	1	0.0472	0.0283	0.0553	0.0197	0.0548
$\lambda/\lambda_\beta = 1/\beta$	%	46.3	18.3	51.3	22.9	47.3
SUMXMY2	Pa ²	1.400	1.399	21.000	1.100	60.902
strain explicit						
$\varepsilon(0)$	1	0	0	2.5	0	4.3
α^a	yr ⁻¹	0.0247	0.0214	0.0257	0.0162	0.0270

^a Given in yr⁻¹.

^b Derived for K and D deviating from their respective mean values equally in relative terms.

Tab. SI10-2: Like Table SI10-1; with the difference that Table SI10-2 comprises the parameters of the three 1959–2015 periods in terms of min-max intervals. The background colors of the cells in Table SI10-1 are preserved.

Parameters		C	B	Min–Max: Case 0 and B and C	
		1850–1958	1900–1958	1959–2015	
stress explicit					
$\sigma(0)$	Pa	0	0	---	---
K	Pa	2.1	2.4	9.9	12.7
D	Pa yr	467.9	467.7	459.2	461.5
$\lambda^{a,b}$	yr ⁻¹	0.0045	0.0051	0.0214	0.0276
λ^{-1}	yr	223.5	196.3	36.3	46.8
α^a	yr ⁻¹	0.0151	0.0228	0.0247	0.0281
β	1	4.371	5.475	1.951	2.158
λ_β^a	yr ⁻¹	0.0196	0.0279	0.0461	0.0538
λ_β^{-1}	yr	51.1	35.9	18.6	21.7
q_β	1	0.9806	0.9725	0.9476	0.9549
T_∞	1	50.6	35.4	18.1	21.2
M_∞ = T_∞/q_β	1	51.6	36.4	19.1	22.2
P_∞ = $1/T_\infty$	1	0.0197	0.0283	0.0472	0.0553
$\lambda/\lambda_\beta = 1/\beta$	%	22.9	18.3	46.3	51.3
SUMXMY2	Pa ²	1.100	1.399	---	---
strain explicit					
$\varepsilon(0)$	1	0	0	---	---
α^a	yr ⁻¹	0.0162	0.0214	0.0247	0.0270

Acronyms and Nomenclature (used in Ms No. esd-2021-27 and in this SI)

ad	adiabatic
C	carbon
comb	combined
CO ₂	carbon dioxide (chemical formula)
CO ₂	atmospheric CO ₂ concentration (in ppmv; parameter)
D	damping constant (in Pa yr)
DIC	dissolved inorganic carbon (in $\mu\text{mol kg}^{-1}$)
E	Young's modulus (in Pa)
GHG	greenhouse gas
h	altitude (in m)
it	isothermal
K	compression modulus (in Pa)
L	land (index)
L	leaf-level factor (in ppmv ⁻¹ ; parameter)
M	memory (in units of 1)
MB	Maxwell body
n.a.	not assessable
NPP	net primary productivity/production (in PgC yr ⁻¹)
O	oceans
p	atmospheric pressure (in hPa)
pCO ₂	partial pressure of atmospheric CO ₂ (in μatm)
P	persistence (in units of 1)
Ph	global photosynthetic carbon influx (in PgC yr ⁻¹)
q	auxiliary quantity (in units of 1)
red	reduced
R	Revelle (buffer) factor (in units of 1)
SD	supplementary data
SE	sensitivity experiment
SI	supplementary information
t	time (in yr)
T	delay time (in units of 1)
TOA	top of the atmosphere
w	weight(ed)
α	exponential growth factor of the strain (in yr ⁻¹)
α_{ppm}	exponential growth factor of the atmospheric CO ₂ concentration (in yr ⁻¹)
β	auxiliary quantity (in units of 1)
β_{b}	biotic growth factor (in units of 1)
β_{Ph}	photosynthetic beta factor (in units of 1)
ε	strain (referring to atmospheric expansion by volume and CO ₂ uptake by sinks; in units of 1)
γ	isentropic coefficient of expansion (in units of 1)

- κ compressibility (in Pa^{-1})
- σ stress (atmospheric CO_2 emissions from fossil fuel burning and land use; in Pa)

References

1. Wikibooks: Formelsammlung Mathematik: Endliche Reihen, Sektion 10: Partialsummen der geometrischen Reihe, https://de.wikibooks.org/wiki/Formelsammlung_Mathematik:_Endliche_Reihen#Partialsummen_der_geometrischen_Reihe (last access 6 February 2022), 24 November 2019.
2. 2 Degrees Institute: Data sources, Sechelt BC, Canada, <https://www.co2levels.org>, April 2016 (global atmospheric concentrations of carbon dioxide over time: https://www.epa.gov/sites/production/files/2016-08/ghg-concentrations_fig-1.csv; last access 6 February 2022).
3. National Oceanic and Atmospheric Administration: Trends in atmospheric carbon dioxide, Silver Springs MD, United States of America, <https://www.esrl.noaa.gov/gmd/ccgg/trends/data.html>, 2022 (Mouna Loa CO₂ annual mean data: ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2_gr_mlo.txt; last access 6 February 2022).
4. Global Carbon Project: Global carbon budget 2019, <https://www.icos-cp.eu/science-and-impact/global-carbon-budget/2019>, 4 December 2019 (data: http://www.globalcarbonproject.org/carbonbudget/archive/2016/Global_Carbon_Budget_2016v1.0.xlsx; see Worksheet *Historical Budget* therein for data prior to 1958, and Worksheet *Fossil Emissions by Fuel Type* and Worksheet *Land-Use Change Emissions* for data from 1959 to 2015; last access 6 February 2006).
5. O'Sullivan, M., Spracklen, D. V., Batterman, S. A., Arnold, S. R., Gloor, M., and Buermann, W.: Have synergies between nitrogen deposition and atmospheric CO₂ driven the recent enhancement of the terrestrial carbon sink? *Global Biogeochem. Cycles*, 33, 163–180, <https://doi.org/10.1029/2018GB005922>, 2019 (research data Leeds repository: <http://archive.researchdata.leeds.ac.uk/482/>; last access: 28 February 2022).
6. Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J., and Santana-Casiano, J. M.: A time-series view of changing ocean chemistry due to ocean uptake of anthropogenic CO₂ and ocean acidification, *Oceanography*, 27, 126–141, <https://doi.org/10.5670/oceanog.2014.16>, 2014 (supporting online material: https://tos.org/oceanography/assets/images/content/27-1_bates_supplement.pdf; last access 6 February 2022).
7. CDIAC: Carbon Dioxide Information Analysis Center - Conversion tables, <https://cdiac.ess-dive.lbl.gov/pns/convert.html> (last access 6 February 2022), 26 September 2012.
8. Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and other biogeochemical cycles, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., Cambridge University Press, Cambridge, United Kingdom, 465–570, https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter09_FINAL.pdf, ISBN: 978-1-107-66182-0, 2013.
9. Silberberg, M. S.: *Chemistry: The Molecular Nature of Matter and Change*, McGraw-Hill, Boston, MA, United States of America, ISBN: 9780077216504, 2009 (background source to https://en.wikipedia.org/wiki/Dalton%27s_law; last access 6 February 2022).
10. Luo, Y., and Mooney, H. A.: Stimulation of global photosynthetic carbon influx by an increase in atmospheric carbon dioxide concentration, in: *Carbon Dioxide and Terrestrial Ecosystems*, edited by Koch, G. W., and Mooney, H. A., Academic Press, San Diego, United States of America, 381–397, ISBN: 0-12-505295-2, 1996.
11. Wark, K.: *Thermodynamics*, McGraw2Hill, New York NY, United States of America, ISBN: 9780070682849, 1983 (background source to http://homepages.wmich.edu/~cho/ME432/Appendix1_SUnits.pdf; cf. also https://en.wikipedia.org/wiki/Heat_capacity_ratio; last access 28 January 2022).
12. Lackner, B. C., Steiner, A. K., Hegerl, G. C., and Kirchengast, G.: Atmospheric climate change detection by radio occultation using a fingerprinting method, *J. Climate*, 24, 5275–5291, <https://doi.org/10.1175/2011JCLI3966.1>, 2011.

13. Steiner, A. K., Lackner, B. C., Ladstädter, F., Scherllin-Pirscher, B., Foelsche, U., and Kirchengast, G.: GPS radio occultation for climate monitoring and change detection, *Radio Sci.*, 46, RS0D24 (17pp), <https://doi.org/10.1029/2010RS004614>, 2011.
14. Cavcar, M.: The international standard atmosphere (ISA), Anadolu University, Eskişehir, Turkey (7pp), <http://fisicaatmo.at.fcen.uba.ar/practicas/ISAweb.pdf> (last access: 28 February 2022), 2000.
15. Wikipedia: List of integrals of exponential functions, https://en.wikipedia.org/wiki/List_of_integrals_of_exponential_functions (last access 6 February 2022), 31 January 2022.