

# ***WORKING PAPER***

## **QUALITATIVE SIMULATION OF DIFFERENTIAL EQUATIONS**

*J.-P. Aubin*

March 1988  
WP-88-015

# QUALITATIVE SIMULATION OF DIFFERENTIAL EQUATIONS

*J.-P. Aubin*

March 1988  
WP-88-015

*Working Papers* are interim reports on work of the International Institute for Applied Systems Analysis and have received only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute or of its National Member Organizations.

INTERNATIONAL INSTITUTE FOR APPLIED SYSTEMS ANALYSIS  
A-2361 Laxenburg, Austria

# **Qualitative Simulation of Differential Equations**

**JEAN-PIERRE AUBIN**

**CEREMADE, UNIVERSITÉ DE PARIS-DAUPHINE  
& IIASA, INTERNATIONAL INSTITUTE FOR APPLIED SYSTEMS  
ANALYSIS**

## **FOREWORD**

This paper deals with the QSIM algorithm introduced by Kuipers to track down the changes of monotonicity properties of solutions to a differential equation or other observations of the solutions. It introduces the concept of “qualitative cells” where the monotonicity properties of the states of the system remain the same. It provides sufficient conditions for the non emptiness of such cells, for their singularities, for the transition from one cell to another, and characterizes also the qualitative equilibria and repellers of the associated qualitative system.

Alexander B. Kurzhanski  
Chairman  
System and Decision Sciences Program

## **Contents**

<b>1</b>	<b>Transitions between qualitative cells</b>	<b>4</b>
<b>2</b>	<b>Qualitative Equilibrium and Repellor</b>	<b>7</b>
<b>3</b>	<b>The QSIM Algorithm</b>	<b>9</b>
<b>4</b>	<b>Nonemptiness and Singularity of Qualitative Cells</b>	<b>11</b>

## Introduction

The purpose of this paper is to revisit the QSIM algorithm introduced by Kuipers in [25] for studying the qualitative evolution of solutions to a differential equation

$$(1) \quad \dot{x}(t) = f(x(t))$$

where the state  $x$  ranges over a closed subset  $K$  of a finite dimensional vector-space  $X := \mathbf{R}^n$ . We posit the assumptions of Nagumo Theorem<sup>1</sup>:

$$(2) \quad \begin{cases} \text{i)} & f \text{ is continuous with linear growth} \\ \text{ii)} & K \text{ is a closed viability domain} \end{cases}$$

The **qualitative state** of a solution to the differential equation (1) at a given time  $t$  is the knowledge of the monotonicity property of each component  $x_i(t)$  of a solution  $x(\cdot)$  to this differential equation, i.e., the knowledge of the sign of the derivatives  $x'_i(t)$ . Hence the **qualitative behavior** is the evolution of the qualitative states of the solution, i.e., the evolution of the vector of signs of the components of  $x'(t) = f(x(t))$ , which must be determined without solving the differential equation.

In order to denote the qualitative states and track down their evolution, we introduce<sup>2</sup> the strict and large confluence frames  $(\mathcal{R}^n, Q_n)$  and  $(\mathcal{R}^n, \bar{Q}_n)$  of  $\mathbf{R}^n$  where  $\mathcal{R}^n$  denotes the  $n$ -dimensional **confluence space** defined by

$$\mathcal{R}^n := \{-, 0, +\}^n$$

where  $Q_n$  is the set-valued map from  $\mathcal{R}^n$  to  $\mathbf{R}^n$  associating with every  $a \in \mathcal{R}^n$  the convex cone

$$Q_n(a) := \mathbf{R}_a^n := \{v \in \mathbf{R}^n \mid \text{sign of } (v_i) = a_i\}$$

---

<sup>1</sup>We recall that the **contingent cone**  $T_K(x)$  to a subset  $K$  at  $x \in K$  is the closed cone of elements  $v$  satisfying

$$\liminf_{h \rightarrow 0^+} d(x + hv, K)/h = 0$$

and that  $K$  is a viability domain if and only if

$$\forall x \in K, f(x) \in T_K(x)$$

**NAGUMO THEOREM** — Under assumptions (2)i), a closed subset  $K$  is a viability domain if and only if enjoys the viability property: for any initial state  $x_0 \in K$ , there exists a solution  $x(\cdot)$  to the differential equation (1), which is **viable** in the sense that  $x(t)$  remains in  $K$  for all  $t \geq 0$ . [1, Chapter 4]

<sup>2</sup>See [2,6]

and  $\bar{Q}_n$  the set-valued map associating with every  $a \in \mathcal{R}^n$  their closure

$$\bar{Q}_n(a) := a\mathbf{R}_+^n := \{v \in \mathbf{R}^n \mid \text{sign of } (v_i) = a_i \text{ or } 0\}$$

We observe that the inverse of the set-valued map  $Q_n$  is the single-valued map  $s_n$  from  $\mathbf{R}^n$  to  $\mathcal{R}^n$  defined by:

$$\forall i \in \{1, \dots, n\}, s_n(x)_i := \text{sign of } x_i$$

For studying the qualitative behavior of the differential equation (1), i.e., the evolution of the functions  $t \mapsto s_n(x'(t))$  associated to solutions  $x(\cdot)$  of the differential equation, we split the viability domain  $K$  of the differential equation into  $3^n$  “qualitative cells”  $K_a$  and “large qualitative cells”  $\bar{K}_a$  defined by

$$(3) \quad K_a := \{x \in K \mid f(x) \in \mathbf{R}_a^n\} \ \& \ \bar{K}_a := \{x \in K \mid f(x) \in a\mathbf{R}^n\}$$

Indeed, the quantitative states  $x(\cdot)$  evolving in a given qualitative cell  $K_a$  share the same monotonicity properties because, as long as  $x(t)$  remains in  $K_a$ ,

$$\forall i = 1, \dots, n, \text{ sign of } \frac{dx_i(t)}{dt} = a_i$$

**The qualitative cell  $K_0$  is then the set of the equilibria<sup>3</sup> of the system, because  $K_0 = \{x \in K \mid f(x) = 0\}$ .**

Studying the qualitative evolution of the differential equation amounts to know the laws (if any) which govern the transition from one qualitative cell  $K_a$  to other cells **without solving the differential equation.**

But before proceeding further, we shall generalize our problem — free of any mathematical cost — to take care of physical considerations.

Instead of studying the monotonicity properties of each component  $x_i(\cdot)$  of the state of the system under investigation, which can be too numerous, we shall only study the monotonicity properties of  $m$  functionals  $V_j(x(\cdot))$  on the state (for instance, energy or entropy functionals in physics, observations in control theory, various economic indexes in economics) which do matter.

The previous case is the particular case when we take the  $n$  functionals  $V_i$  defined by  $V_i(x) := x_i$ .

We shall assume for simplicity that these functionals  $V_j$  are continuously differentiable around the viability domain  $K$ .

---

<sup>3</sup>Such an equilibrium does exist whenever the viability domain  $K$  is convex and compact. (see [3, Theorem 6.4.11, p. 341]).

We denote by  $\mathbf{V}$  the map from  $X$  to  $Y := \mathbf{R}^m$  defined by

$$\mathbf{V}(x) := (V_1(x), \dots, V_m(x))$$

and we introduce the strict and large confluence frames  $(\mathcal{R}^m, Q_m)$  and  $(\mathcal{R}^m, \overline{Q}_m)$  of  $Y$  for studying the qualitative evolution of the observation  $\mathbf{V}(x(\cdot))$ .

Since the derivative of the observation  $\mathbf{V}(x(\cdot))$  is equal to  $\mathbf{V}'(x(\cdot))x'(\cdot) = \mathbf{V}'(x(\cdot))f(x(\cdot))$ , it will be convenient to set

$$(4) \quad \forall x \in K, \quad g(x) := \mathbf{V}'(x)f(x)$$

(We observe that  $g$  is equal to  $f$  in the particular case).

Hence, we associate with each qualitative state  $a$  the qualitative cells  $K_a$  and the large qualitative cells  $\overline{K}_a$  defined by

$$(5) \quad K_a := \{x \in K \mid g(x) \in \mathbf{R}_a^m\} \ \& \ \overline{K}_a := \{x \in K \mid g(x) \in a\mathbf{R}^m\}$$

In other words, the quantitative states  $x(\cdot)$  evolving in a given qualitative cell  $K_a$  share the same monotonicity properties of their observations because, as long as  $x(t)$  remains in  $K_a$ ,

$$\forall j = 1, \dots, m, \quad \text{sign of } \frac{d}{dt}V_j(x(t)) = a_j$$

In particular, the  $m$  functions  $V_j(x(t))$  remain constant while they evolve in the qualitative cell  $K_0$ .

By using observation functionals chosen in such a way that many qualitative cells are empty, the study of transitions may be drastically simplified: this is a second reason to carry our study in this more general setting.

This the case for instance when the observation functionals are "Lyapunov functions"  $V_j : K \mapsto \mathbf{R}$ . We recall that  $V$  is a Lyapunov function if  $\langle V'(x), f(x) \rangle \leq 0$  for all  $x \in K$ , so that  $V(x(\cdot))$  decreases along the solutions to the differential equation.

Hence, if the observation functionals are Lyapunov functions, the qualitative cells  $K_a$  are empty whenever a component  $a_i$  is positive. In this case, we have at most  $2^m$  non empty qualitative cells. (In some sense, one can say that Lyapunov was the originator of qualitative simulation a century ago).

Here again, studying the qualitative evolution of the differential equation amounts to find the laws which govern the transition from one qualitative cell  $K_a$  to other qualitative cells.



Naturally, we would like to know this evolution directly without solving the differential equation, and therefore, without knowing the state of the system, but only some of its properties.

In other words, the problem arises whether we can map the differential equation (1) to a discrete dynamical system  $\Phi : \mathcal{R}^m \rightsquigarrow \mathcal{R}^m$  on the qualitative space  $\mathcal{R}^m$ .

This is not always possible, and we have thus to define the class of differential equations which enjoy this property.

But before, we shall characterize the qualitative equilibria, which are the qualitative states  $a$  such that the solutions which arrive in the qualitative cell  $\overline{K}_a$  remain in this cell, as well as the qualitative repellers  $b$ , such that any solution which arrive in  $\overline{K}_b$  must leave this cell in finite time.

We shall finally provide conditions insuring that the qualitative cells are not empty or singular around  $\bar{x} \in K_0$  in the sense that there is no other  $x \in \overline{K}_a$  in the neighborhood of  $\bar{x}$ .

## 1 Transitions between qualitative cells

We shall assume from now on that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ .

Let us denote by  $S : K \mapsto \mathcal{C}^1(0, \infty; X)$  the “solution map” associating with each initial state  $x_0 \in K$  the solution  $Sx_0(\cdot)$  to the differential equation (1) starting at  $x_0$  (which is unique since  $f$  is locally lipschitzean).

**Definition 1.1** *Let us consider a map  $f$  from  $K$  to  $X$  and  $m$  observation functionals  $V_j : K \mapsto \mathbf{R}$ . We denote by  $\mathcal{D}(f, \mathbf{V})$ , the “qualitative domain of  $(f, \mathbf{V})$ ”, the subset of qualitative states  $a \in \mathcal{R}^n$  such that the associated qualitative cell  $K_a$  is not empty.*

*We shall say that a qualitative state  $c \in \mathcal{D}(f, \mathbf{V})$  is a “successor” of a qualitative state  $b \in \mathcal{D}(f, \mathbf{V})$  if for all initial state  $x_0 \in \overline{K}_b \cap \overline{K}_c$ , there exists  $\tau \in ]0, +\infty[$  such that  $Sx_0(s) \in K_c$  for all  $s \in ]0, \tau[$ .*

*A qualitative state  $a \in \mathcal{D}(f, \mathbf{V})$  is said to be a “qualitative equilibrium” if it is its own successor. It is said to be a “qualitative repeller” if for any initial state  $x_0 \in \overline{K}_a$ , there exists  $t > 0$  such that  $Sx_0(t) \notin \overline{K}_a$ .*

Our first objective is to express the fact that  $c$  is a successor of  $b$  through a set-valued map  $\Phi$  that we shall define.

For that purpose, we shall set

$$h(x) := g'(x)f(x) = V''(x)(f(x), f(x)) + V'(x)f'(x)f(x)$$

because  $g'(x)v = V''(x)(f(x), v) + V'(x)f'(x)v$ .

We introduce the notation

$$\overline{K}_a^i := \{ x \in \overline{K}_a \mid g(x)_i = 0 \}$$

(Naturally,  $\overline{K}_a = \overline{K}_a^i$  whenever  $a_i = 0$ .)

We shall denote by  $\Gamma$  the set-valued map from  $\mathcal{R}^m$  to itself defined by

(6)  $\forall a \in \mathcal{R}^m$ ,  $(\Gamma(a))_i$  is the set of signs of  $h_i(x) = g'(x)f(x)_i$  when  $x \in \overline{K}_a^i$

We also set  $I_0(x) := \{ i = 1, \dots, m \mid g(x)_i = 0 \}$  and

$$\mathbf{R}_+^{I_0(x)} := \{ v \in \mathbf{R}^m \mid v_i \geq 0 \ \forall i \in I_0(x) \}$$

We introduce the operations  $\wedge$  on  $\mathcal{R}^m$  defined by

$$(b \wedge c)_i := \begin{cases} b_i & \text{if } b_i = c_i \\ 0 & \text{if } b_i \neq c_i \end{cases}$$

and the set-valued operation  $\vee$  where  $b \vee c$  is the subset of qualitative states  $a$  such that

$$a_i := b_i \text{ or } c_i$$

We set

$$a \# b \iff \forall i = 1, \dots, m, a_i \neq b_i$$

**Proposition 1.1** *The set-valued map  $\Gamma$  satisfies the consistency property*

$$(7) \quad \Gamma(a \vee 0) \subset \Gamma(a)$$

and thus,

$$\Gamma(b \wedge c) \subset \Gamma(b) \cap \Gamma(c)$$

**Proof** — To say that  $\overline{K}_b$  is contained in  $\overline{K}_a$  amounts to saying that  $b$  belongs to  $a \vee 0$ . When this is the case, we deduce that for all  $i = 1, \dots, m$ ,  $\overline{K}_b^i \subset \overline{K}_a^i$ , so that the signs taken by  $h(x)_i$  when  $x$  ranges over  $\overline{K}_b^i$  belong to the set of  $\Gamma(a)_i$  of signs taken by the same function over  $\overline{K}_a^i$ . Therefore,  $\Gamma(b)$  is contained in  $\Gamma(a)$ .

Since  $b \wedge c$  belongs to both  $b \wedge 0$  and  $c \vee 0$ , we deduce that  $\Gamma(b \wedge c)$  is contained in both  $\Gamma(b)$  and  $\Gamma(c)$ .  $\square$

**Definition 1.2** We shall associate with the system  $(f, \mathbf{V})$  the discrete dynamical system on the confluence set  $\mathcal{R}^m$  defined by the set-valued map  $\Phi : \mathcal{R}^m \rightsquigarrow \mathcal{R}^m$  associating with any qualitative state  $b$  the subset

$$(8) \quad \Phi(b) := \{ c \in \mathcal{D}(f, \mathbf{V}) \mid \Gamma(b \wedge c) \subset c \vee 0$$

We begin with necessary conditions for a qualitative state  $c$  to be a successor of  $b$ :

**Proposition 1.2** Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ .

If  $c \in \mathcal{D}(f, \mathbf{V})$  is a successor of  $b$ , then  $c$  belongs to  $\Phi(b)$ .

Before proving this proposition, we need the following

**Lemma 1.1** Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ .

If  $v$  belongs to the contingent cone to the large qualitative cell  $\overline{K}_a$ , then the condition

$$(9) \quad v \in T_K(x) \ \& \ \forall i \in I_0(x), \ \text{sign of } (g'(x)v)_i = a_i \ \text{or } 0$$

is satisfied.

The converse is true if we posit the transversality assumption

$$\forall x \in \overline{K}_a, \ g'(x)C_K(x) - a\mathbf{R}_+^{I_0(x)} = \mathbf{R}^m$$

**Proof** — Since the large qualitative cell  $\overline{K}_a$  is the intersection of  $K$  with the inverse image by  $g$  of the convex cone  $a\mathbf{R}_+^m$ , we know<sup>4</sup> the contingent cone to  $\overline{K}_a$  at some  $x \in \overline{K}_a$  is contained into

$$T_{\overline{K}_a}(x) = T_K(x) \cap g'(x)^{-1}T_{a\mathbf{R}_+^m}(g(x))$$

and is equal to this intersection provided that the “transversality assumption”

$$g'(x)C_K(x) - C_{a\mathbf{R}_+^m}(g(x)) = \mathbf{R}^m$$

is satisfied. On the other hand, we know that the  $a\mathbf{R}_+^m$  cone being convex,

$$C_{a\mathbf{R}_+^m}(y) = T_{a\mathbf{R}_+^m}(y) = aT_{\mathbf{R}_+^m}(ay) \supset a\mathbf{R}_+^m$$

---

<sup>4</sup>See [3, Theorem 7.3. p.]

and that  $v \in T_{\mathbf{R}_+^m}(z)$  if and only if

$$\text{whenever } z_j = 0, \text{ then } v_j \geq 0$$

Consequently,  $v \in T_{\mathbf{aR}_+^m}(g(x))$  if and only if

$$\text{whenever } g(x)_j = 0, \text{ then sign of } v_j = a_j \text{ or } 0$$

i.e.,  $T_{\mathbf{aR}_+^m}(g(x)) = \mathbf{aR}_+^{I_0(x)}$ .

Hence  $v$  belongs to the contingent cone to  $\overline{K_a}$  at  $x$  if and only if  $v$  belongs to  $T_K(x)$  and  $g'(x)v$  belongs to  $T_{\mathbf{aR}_+^m}(g(x))$ , i.e., the sign of  $(g'(x)v)_j$  is equal to  $a_j$  or 0 whenever  $j$  belongs to  $I_0(x)$ .  $\square$

**Proof of Proposition 1.2** — Let  $c$  be a successor of  $b$ . Take any initial state  $x_0$  in  $\overline{K_b} \cap \overline{K_c}$  and set  $x(t) := S_{x_0}(t)$ . We observe that the intersection of two qualitative cells  $\overline{K_b}$  and  $\overline{K_c}$  is equal to

$$\overline{K_b} \cap \overline{K_c} := \overline{K_{b \wedge c}}$$

Since the solution  $x(t)$  to the differential equation crosses the intersection  $\overline{K_{b \wedge c}}$  towards  $\overline{K_c}$ ,  $f(x_0)$  belongs to the contingent cone  $T_{\overline{K_c}}(x_0)$  because

$$\liminf_{h \rightarrow 0^+} d_{K_c}(x_0 + hf(x_0))/h \leq \liminf_{h \rightarrow 0^+} \left\| x'(0) - \frac{x(h) - x_0}{h} \right\| = 0$$

By Lemma 1.1, this implies that

$$\forall x_0 \in \overline{K_{b \wedge c}}, \forall i \in I_0(x_0), \text{ sign of } g'(x_0)f(x_0)_i = c_i \text{ or } 0$$

or, equivalently, that

$$\Gamma(b \wedge c) \subset c \vee 0$$

Hence  $c$  belongs to  $\Phi(b)$ , as it was announced.  $\square$

## 2 Qualitative Equilibrium and Repellor

We can characterize the qualitative equilibria of the differential equation.

**Theorem 2.1** *Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ . We posit the transversality assumption*

$$(10) \quad \forall x \in \overline{K_a}, g'(x)C_K(x) - \mathbf{aR}_+^{I_0(x)} = \mathbf{R}^m$$

*Then  $a$  is a qualitative equilibrium if and only if  $a$  belongs to  $\Phi(a)$ .*

**Proof** — We already know that if  $a$  is a qualitative equilibrium, then  $a$  belongs to  $\Phi(a)$ . We shall prove the converse statement, and, for that purpose, observe that saying that  $a$  is a qualitative equilibrium amounts to saying that the large qualitative cell  $\overline{K}_a$  enjoys the viability property (or is invariant by  $f$ ). By the Nagumo Theorem, this is equivalent to say that  $\overline{K}_a$  is a viability domain, i.e., that

$$\forall x \in \overline{K}_a, f(x) \in T_{\overline{K}_a}(x)$$

By the Lemma 1.1, knowing that  $f(x)$  belongs to the contingent cone  $T_K(x)$  by assumption, this amounts to say that

$$\forall x \in \overline{K}_a, \forall i \in I_0(x), \text{ sign of } (g'(x)f(x))_i = a_i \text{ or } 0$$

i.e., that  $\Gamma(a \wedge a) = \Gamma(a) \subset a \vee 0$ . Hence,  $a$  is a fixed point of  $\Phi$ .  $\square$

What happens if a large qualitative cell  $\overline{K}_a$  is not a viability domain of  $f$ ? We know<sup>5</sup> that there exists a closed largest viability domain of  $f$  contained in  $K_a$ , called the viability kernel  $\text{Viab}(\overline{K}_a)$ .

We infer from the definition of the viability kernel that

**Proposition 2.1** *Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ . We posit the transversality assumption*

$$(11) \quad \forall x \in \overline{K}_a, g'(x)C_K(x) - a\mathbf{R}_+^{I_0(x)} = \mathbf{R}^m$$

— *The qualitative state  $a$  is a qualitative repellor if and only if the viability kernel of  $\overline{K}_a$  is empty.*

— *If for some  $b \in a \vee 0$ , the qualitative cell  $\overline{K}_b$  is contained in the viability kernel  $\text{Viab}(\overline{K}_a)$ , then  $a$  is the only successor of  $b$ .*

**Proof**

1 — To say that some  $x_0 \in \overline{K}_a$  does not belong to the viability kernel of  $\overline{K}_a$  means that for some  $t > 0$ ,  $Sx_0(t) \notin \overline{K}_a$ . If this happens for all  $x_0 \in \overline{K}_a$ , then obviously,  $a$  is a qualitative repellor.

2 — If  $\overline{K}_b \subset \text{Viab}(\overline{K}_a)$ , then, for all  $x_0 \in \overline{K}_b$ ,  $Sx_0(t) \in \overline{K}_a$  for all  $t \geq 0$ . Hence  $a$  is the only successor of  $b$ .  $\square$

<sup>5</sup>The viability kernel  $\text{Viab}(M)$  of a closed subset  $M \subset K$  of a viability domain  $K$  of a  $f$  is the largest closed viability domain of the restriction  $f|_M$  of  $f$ . It is the subset of elements  $x \in M$  such that  $Sx(t) \in M$  for all  $t \geq 0$ . It may naturally be empty.

### 3 The QSIM Algorithm

We shall now distinguish the  $2^n$  “full qualitative states”  $a \neq 0$  from the other qualitative states, the “transition states”.

When  $I$  is a non empty subset of  $N := \{1, \dots, m\}$ , we associate with a full state  $a \neq 0$  the transition state  $a^I$  defined by

$$a_i^I := \begin{cases} 0 & \text{if } i \in I \\ a_i & \text{if } i \notin I \end{cases}$$

**Lemma 3.1** *Let  $a \neq 0$  be qualitative state which is not a qualitative equilibrium. There exist a solution starting at some  $x \in K_a$  and some  $t_1 > 0$  such that  $x(t) \in K_a$  for  $t \in [0, t_1[$  and  $x(t_1) \in K_{a^I}$  for some non empty subset  $I \subset N$ :*

$$\forall i \in I, x(t_1) \in K_a^i$$

**Proof** — Let us choose  $x \in K_a$  and set  $x(t) := Sx(t)$ ,  $x(0) = x$  for simplicity. Either  $x(t)$  remains in  $\overline{K}_a$  for all  $t$ , or there exists  $\tau > 0$  such that  $x(\tau) \notin \overline{K}_a$ . Since  $a$  is not a qualitative equilibrium, the latter happens for at least one initial state  $x$ .

Let  $J_a := \{t > 0 \mid x(t) \notin \overline{K}_a\}$ , an open subset of  $\mathbf{R}_+$  and  $t_1 := \inf J_a$ .

Since  $a \neq 0$  is a full cell and since the initial point  $x$  belongs to the cell  $\mathbf{R}_a^m$ , which is open, then  $x(t)$  remains in  $\mathbf{R}_a^m$  for  $t \in ]0, \eta[$  for some  $\eta > 0$ , and thus,  $t_1 > 0$ . Since  $x(t) \in K_a$  for all  $t < t_1$ , we deduce that  $x(t_1)$  belongs to a transition cell  $\overline{K}_a$ .

By definition of  $t_1$ , there exists a sequence  $t_p > t_1$  converging to  $t_1$  such that  $x(t_p) \notin \overline{K}_a$ , i.e., such that  $x(t_p) \in K$  and  $g(x(t_p)) \notin a\mathbf{R}_+^n$ . This means that there exists a non empty subset  $I \subset N$  such that  $g(x(t_1))_i = 0$  for all  $i \in I$ .  $\square$

We face now two types of problems:

1 — What are the transition states  $a^I \in a \vee 0$  such that the cell  $K_{a^I}$  is reached in finite time by at least a solution starting in  $K_a$ ?

This problem is closely related to the “target problem” and other controllability issues in control theory, which received only partial solutions. We shall not attempt to answer this question in this paper.

2 — What are the successors, if any, of a given transition state  $a^I$ ?

The second question does not always receive an answer, since, starting from some initial state  $x \in K_a^I$ , there may exist two sequences  $t_n > 0$  and  $s_n > 0$  converging to  $0+$  such that  $x(t_n) \in \overline{K}_a$  and  $x(s_n) \notin \overline{K}_a$ .

We can exclude this pathological phenomenon in two instances.

One obviously happens when either  $a$  or the transition state  $a^I$  is an equilibrium, i.e., when

$$\Gamma(a)_i = \begin{cases} 0 & \text{when } i \in I \\ \in \{a_i, 0\} & \text{when } i \notin I \end{cases}$$

This also happens in the following situation:

**Lemma 3.2** *Let  $a \neq 0$  be a full transition state. If  $\Gamma(a) \neq 0$  (and thus, is reduced to a point) then, for all transition state  $a^I$ , there exists a unique successor  $b := \Phi(a^I) \neq 0$  : for all initial state  $x$  in the transition cell  $K_{a^I}$ , there exists  $t_2 > 0$  such that, for all  $t \in ]0, t_2[$ , the solution  $x(t)$  remains in the full qualitative cell  $K_b$ .*

**Proof** — We consider an initial state  $x \in K_{a^I}$ .

If  $i \notin I$ , then the sign of  $g(x)_i$  is equal to  $a_i \neq 0$ , and thus, there exists  $\eta_i > 0$  such that the sign of  $g(x(t))_i$  remains equal to  $a_i^I = a_i$  when  $t \in ]0, \eta_i[$ .

If  $i \in I$ , then  $g(x)_i = 0$ , and we know that the sign of the derivative  $\frac{d}{dt}g_i(x(t))|_{t=0} = h_i(x)$  is equal to  $b_i := \Gamma(a)_i$  and is different from 0. Hence there exists  $\eta_i > 0$  such that the sign of  $h(x(t))_i$  remains equal to  $b_i$  when  $t \in ]0, \eta_i[$ , so that the sign of

$$g_i(x(t)) = \int_{t_1}^t h_i(x(\tau)) d\tau$$

remains equal to  $\Gamma(a)_i$  on the interval  $]0, \eta_i[$ .

Hence we have proved that there exists some  $\eta > 0$  such that  $x(t) \in K_b$  for  $t \in ]0, \eta[$  where

$$b_i := \begin{cases} \Gamma(a)_i & \text{when } i \in I \\ a_i & \text{when } i \notin I \end{cases}$$

and where  $t_2 := \min_{i \in N} \eta_i > 0$ .  $\square$

**Definition 3.1** *We shall say that the system  $(f, V_j)$  is “strictly filterable” if and only if for all full state  $a \in D(f, \mathbf{V}) \neq 0$ , either  $\Gamma(a) \neq 0$  or  $a$  is a qualitative equilibrium or all the transition states  $a^I$  ( $I \neq \emptyset$ ) are qualitative equilibria.*

We deduce from Definition 3.1 and the above observations the following consequence:

**Theorem 3.1** *Let us assume that  $f$  is continuously differentiable, that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$  and that the system  $(f, V_j)$  is “strictly filterable”. Let  $a \in \mathcal{R}^m$  be an initial full qualitative state.*

*Then, for any initial state  $x \in K_a$  in the qualitative cell  $K_a$ , the sign vector  $a_x(t) := s_n(Sx(t))$  is a solution to the QSIM algorithm defined in the following way:*

*There exist a sequence of qualitative states  $a_k$  satisfying*

$$(12) \quad a_0 := a \ \& \ a_{k+1} \in \Phi(a_k \vee 0)$$

*and a sequence  $t_0 := 0 < t_1 < \dots < t_n < \dots$  such that*

$$(13) \quad \begin{cases} \forall t \in ]t_k, t_{k+1}[, \ a(t) = a_k \\ a(t_{k+1}) = a_k \wedge a_{k+1} \end{cases}$$

In other words, we know that the vector signs of the variations of the observations of the solutions to the differential equation (1) evolve according the set-valued dynamical system (12) and stop when  $a_k$  is either a qualitative equilibrium or all its transition states  $a_k^I$  are qualitative equilibria.

**Remark** — The solutions to the QSIM algorithm (12) do not necessarily represent the evolution of the variations of the vector signs of a solution to the differential equation.

Further studies must bring answers allowing to delete impossible transitions from one full qualitative cell  $\overline{K}_a$  to some of its transition cells  $K^{a^I}$ .

This is the case of a qualitative equilibrium, for instance, since  $a$  is the only successor of itself.  $\square$

Therefore, the QSIM algorithm requires the definition of the set-valued map  $\Gamma : \mathcal{R}^m \rightsquigarrow \mathcal{R}^m$  by computing the signs of the  $m$  functions  $h_j(\cdot)$  on the qualitative cells  $K_a^i$  for all  $i \in N$  and  $a \in \mathcal{D}(f, \mathbf{V}) \neq \emptyset$ .

If by doing so, we observe that the system is strictly filterable, then we know that the set-valued dynamical system (12) contains the evolutions of the vector signs of the  $m$  observations of solutions to the differential equation (1).

## 4 Nonemptiness and Singularity of Qualitative Cells

The question we answer now is whether these qualitative cells are non empty.



**Theorem 4.1** *Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ . Let  $\bar{x}$  belong to the qualitative cell  $K_0$ . We posit the transversality condition:*

$$(14) \quad g'(\bar{x})C_K(\bar{x}) - a\mathbf{R}_+^m = \mathbf{R}^m$$

*Then the qualitative cell  $K_a$  is nonempty and  $\bar{x}$  belongs to its closure. In particular, if*

$$g'(\bar{x})C_K(\bar{x}) = \mathbf{R}^m$$

*then the  $3^m$  qualitative cells  $K_a$  are nonempty. (We have a chaotic situation, since every qualitative behavior can be implemented as a initial qualitative state.)*

**Proof** — We apply the Constrained Inverse Function Theorem<sup>6</sup> of [3,4] to the map  $(x, y) \mapsto g(x) - y$  from  $X \times Y$  to  $Y$  restricted to the closed subset  $K \times a\mathbf{R}_+^m$  at the point  $(\bar{x}, 0)$ . Its Clarke tangent cone is equal to the product  $C_K(\bar{x}) \times a\mathbf{R}_+^m$  of the Clarke tangent cones  $C_K(\bar{x})$  and

$$C_{a\mathbf{R}_+^m}(0) = a\mathbf{R}_+^m$$

Therefore, we know that there exists  $\epsilon > 0$  such that, for all  $z \in \epsilon[-1, +1]^m$ , there exist an element  $x \in K$  and an element  $y \in a\mathbf{R}_+^m$  satisfying  $g(x) - y = z$  and  $\|x - \bar{x}\| + \|y\| \leq l\|z\|$ . Taking in particular  $z_i = a_i\epsilon$ , we see that  $g(x)_i = a_i\epsilon + y_i$  and thus, that the sign of  $g(x)_i$  is equal to  $a_i$  for all  $i = 1, \dots, m$ . Hence  $x$  belongs to  $K_a$  and  $\|x - \bar{x}\| \leq l\epsilon$ .  $\square$

Let  $\bar{x}$  belong to  $K_0$ . We shall say that the qualitative cell  $\bar{K}_a$  is “singular” at  $\bar{x}$  if there exists a neighborhood  $N(\bar{x})$  of  $\bar{x}$  such that  $\bar{x}$  is locally the only point of the qualitative cell  $\bar{K}_a$ :

$$\forall x \in N(\bar{x}) \cap K, \quad x \neq \bar{x}, \quad g(\bar{x}) \notin a\mathbf{R}_+^m$$

**Theorem 4.2** *Let us assume that  $f$  is continuously differentiable and that the  $m$  functions  $V_j$  are twice continuously differentiable around the viability domain  $K$ . Let  $\bar{x}$  belong to the qualitative cell  $K_0$ .*

<sup>6</sup>CONSTRAINED INVERSE FUNCTION THEOREM — Let  $X$  is a Banach space,  $Y$  a finite dimensional vector-space and  $f$  a continuous map from a neighborhood of  $K$  to  $Y$ . Assume that it is continuously differentiable around some point  $x_0 \in K$  and that

$$f'(x_0)C_K(x_0) = Y$$

Then there exists a constant  $l > 0$  such that, for all  $y \in Y$  close enough to  $f(x_0)$ , there exists a solution  $x \in K$  to the equation  $f(x) = y$  satisfying  $\|x - x_0\| \leq l\|y - f(x_0)\|$ .

We posit the following assumption:

$$(15) \quad T_K(\bar{x}) \cap (g'(\bar{x}))^{-1}(a\mathbf{R}_+^m) = \emptyset$$

Then the qualitative cell  $\bar{K}_a$  is singular at  $\bar{x}$ .

**Proof** — We follow the same arguments than in [4]. Assume the contrary: for all  $n > 0$ , there exists  $x_n \in K$ ,  $x_n \neq \bar{x}$  such that  $g(x_n)$  does belong to  $a\mathbf{R}_+^m$ . Let us set  $h_n := \|x_n - \bar{x}\| > 0$  converge to 0 and  $v_n := \left\| \frac{x_n - \bar{x}}{h_n} \right\|$ . Since  $v_n$  belongs to the unit ball, which is compact, a subsequence (again denoted)  $v_n$  converges to some element  $v$  of the unit ball. This limit  $v$  belongs also to the contingent cone  $T_K(\bar{x})$  because, for all  $n > 0$ ,  $\bar{x} + h_n v_n = x_n$  belongs to  $K$ .

Finally, since  $g(\bar{x} + h_n v_n) = g(x_n) \in a\mathbf{R}_+^m$  for all  $n > 0$  and  $g(\bar{x}) = 0$ , we infer that the limit  $g'(\bar{x})v$  of the difference quotients  $\frac{g(\bar{x} + h_n v_n) - g(\bar{x})}{h_n} \in a\mathbf{R}_+^m$  belongs to  $a\mathbf{R}_+^m$ . Hence we have proved the existence of a non zero element

$$v \in T_K(\bar{x}) \cap (g'(\bar{x}))^{-1}(a\mathbf{R}_+^m)$$

a contradiction of the assumption.  $\square$

## References

- [1] AUBIN J.-P. & CELLINA A. (1984) DIFFERENTIAL INCLUSIONS. Springer-Verlag (Grundlehren der Math. Wissenschaften, Vol.264, 1-342)
- [2] AUBIN J.-P. & DORDAN O. (to appear) *Qualitative Frames and Qualitative Projections of Maps.*
- [3] AUBIN J.-P. & EKELAND I. (1984) APPLIED NONLINEAR ANALYSIS. Wiley-Interscience
- [4] AUBIN J.-P. & FRANKOWSKA H. (1987) *On the inverse function Theorem.* J. Math. Pures Appliquées, 66, 71-89
- [5] AUBIN J.-P. (1987) *Differential Calculus of set-valued maps: an update.* IIASA WP
- [6] AUBIN J.-P. (to appear) *Qualitative Equations: the Confluence Case.*
- [7] AUBIN J.-P. (to appear) *Qualitative Simulation of Differential Equations.*
- [8] BOBBROW D. G. (Ed.) (1984) QUALITATIVE REASONING ABOUT PHYSICAL SYSTEMS. Artificial Intelligence Volume 24 (North Holland)
- [9] BOURGINE P. & RAIMAN O. (1986) *Economic as Reasoning on a Qualitative Model.*
- [10] BOURGINE P. & RAIMAN O. (1986) *Macroeconomics as reasoning on a qualitative model.*
- [11] BROWN A.L. (1976) *Qualitative knowledge, causal reasoning and the localization of failures.*
- [12] DAVIS R. (1984) *Diagnostic reasoning based on structure and behavior.* Artificial Intelligence, 24, 347-410
- [13] DE KLEER J. & BOBROW D.J (1984) *Qualitative Reasoning With Higher Order Derivatives.* Artificial Intelligence, 24

- [14] DE KLEER J. & BROWN J.S. (1984) *A Qualitative Physics Based on Confluences*. Artificial Intelligence, 24
- [15] DE KLEER J. & BROWN J.S (1986) *Theories of Causal Ordering* . Artificial Intelligence, 29
- [16] De KLEER J. (1984) *How Circuits Work*. Artificial Intelligence, 24, 205-280
- [17] DE KLEER J. (1986)
- [18] Di MANZO M. & TRUCCO E. (1987) *Commonsense reasoning about flexible objects : A case study*. Advances in Artificial Intelligence, North-Holland
- [19] FORBUS K.D & GENTNER (to appear) *Causal Reasoning about Quantities*. University of Illinois at Urbana-Champaign
- [20] FORBUS K. D. (1984) *Qualitative Process Theory*. Artificial Intelligence, 24, 85-168
- [21] FORBUS K.D. (1987) *The qualitative Process Engine : A Study in Assumption-based Truth Maintenance*. Workshop on Qualitative Physics, Urbana-Champaign ILL
- [22] KUIPERS B. & KASSIRER J.P. (1984) *Causal Reasoning in Medicine : Analysis of a Protocol*. Cognitive Science, 8, 363-385
- [23] KUIPERS B. (1984) *Commonsense Reasoning about Causality: Deriving Behavior from Structure*. Artificial Intelligence, 24, 169-203
- [24] KUIPERS B. (1985) *The Limits of Qualitative Simulation*. Ninth International Joint Conference on Artificial Intelligence, IJCAI-85, Los Angeles
- [25] KUIPERS B. (1986) *Qualitative Simulation*. Artificial Intelligence, 29, 289-338
- [26] KUIPERS B. (1987) *A Reply to Peter Struss*. Draft
- [27] KUIPERS B. (1987) *Abstraction by Time-Scale in Qualitative Simulation*. AAAI, The Sixth National Conference on Artificial Intelligence

- [28] KUIPERS B. (1987) *Qualitative Simulation as Causal Explanation*. IEEE Transaction on Systems, Man, and cybernetics, 17
- [29] LEPETIT M. (1986) *Physique qualitative : le système PHYSICAL*. LRI univèrité de Paris sud
- [30] RAIMAN O. (1985) *Raisonnement Qualitatif*.
- [31] STRUSS P. (1987) *Mathematical Aspects of Qualitative Reasoning*. Workshop on Qualitative Physics, Urbana-Champaign ILL
- [32] TRAVE L. & DORMOY J.L. (1987) *Qualitative Calculus with Applications to Control Theory & Qualitative Physics Problems*. Draft
- [33] TRAVE L. & KASKUREWICZ E. (1986) *Qualitative Controllability & Observability of Linear Dynamical Systems*. IFAC-LSS, Zürich