

# APPENDIX IV

## TEMPERATURE REGIME REQUIREMENTS OF CROPLUTs

### Notes:

**Climates:** B = Boreal; Te = Temperate; STR = Sub-tropics; WR = Winter Rainfall;  
SR = Summer Rainfall; Tr = Tropics

**Growth cycle:** L/Gc = Total; La = Pre-dormancy; Lb = Post-dormancy

### Temperature profile interval symbols:

Temperature intervals (°C)	<-5	-5-0	0-5	5-10	10-15	15-20	20-25	25-30	>30
<b>Totals</b>	L9	L8	L7	L6	L5	L4	L3	L2	L1
<b>Increasing temperatures (winter to summer)</b>	L9a	L8a	L7a	L6a	L5a	L4a	L3a	L2a	L1a
<b>Decreasing temperatures (summer to winter)</b>	L9b	L8b	L7b	L6b	L5b	L4b	L3b	L2b	L1b

**Heat Units:** TSgc = Temperature Sum during growth cycle

**Temperature Growing Period:** LGP<sub>t=5</sub> = Number of days with mean daily temperatures above 5°C

## TEMPERATURE REGIME REQUIREMENTS OF CROPLUTs

Crop	Sub-optimal Conditions	Optimal Conditions
<b>Winter Wheat</b>	<i>Climates: B, Te, STr (WR+SR)</i>	<i>Climates: B, Te, STr (WR+SR)</i>
(Wheat I)	$L6a < 0.667 \cdot Lb$	$L6a < 0.500 \cdot Lb$
<b>Winter Barley</b>	$L6a + L5a > 0.167 \cdot Lb$	$L6a + L5a > 0.084 \cdot Lb$
(Barley I)	$L2a + L2b < 0.333 \cdot Lb$	$L2a + L2b < 0.333 \cdot Lb$
<b>Winter Rye</b>	$L1a + L1b = 0$	$L1a + L1b = 0$
(Rye I)	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$
$L = 30 + 90$	$L3b + L4b + L5b + L6b > La$	$L3b + L4b + L5b + L6b > La$
$L = 35 + 105$	$TSgc > 1200$ *	$TSgc > 1300$ *
$L = 40 + 120$	no permafrost	no permafrost
$L = 45 + 135$	$LGP_{t=5} < 365$	$LGP_{t=5} < 365$
	dormancy required	dormancy required
<b>Spring Wheat</b>	<i>Climates: B, Te, STr (WR+SR)</i>	<i>Climates: B, Te, STr (WR+SR)</i>
(Wheat II)	$L6a < 0.333 \cdot L$	$L6a < 0.333 \cdot L$
$L = 105/120/135/150$	$L6a + L5a > 0.167 \cdot Lb$	$L6a + L5a > 0.167 \cdot Lb$
	$L2a + L2b < 0.333 \cdot Lb$	$L2a + L2b < 0.333 \cdot Lb$
	$L1a + L1b = 0$	$L1a + L1b = 0$
	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$
	$TSgc > 1400$	$TSgc > 1800$
	no permafrost	no permafrost
	$LGP_{t=5} < 365$	$LGP_{t=5} < 365$
<b>Spring Barley</b>	<i>Climates: B, Te, STr (WR+SR)</i>	<i>Climates: B, Te, STr (WR+SR)</i>
(Barley II)	$L6a < 0.333 \cdot L$	$L6a < 0.333 \cdot L$
<b>Spring Rye</b>	$L6a + L5a > 0.167 \cdot Lb$	$L6a + L5a > 0.167 \cdot Lb$
(Rye II)	$L2a + L2b < 0.333 \cdot Lb$	$L2a + L2b < 0.333 \cdot Lb$
$L = 90/105/120/135$	$L1a + L1b = 0$	$L1a + L1b = 0$
	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$	$L2b + L3b + L4b + L5b < 0.500 \cdot Lb$
	$TSgc > 1400$	$TSgc > 1700$
	no permafrost	no permafrost
	$LGP_{t=5} < 365$	$LGP_{t=5} < 365$
<b>Wheat (sub-tropics)</b>	<i>Climates: STr (WR + SR)</i>	<i>Climates: STr (WR + SR)</i>
(Wheat III)	$L6a < 0.333 \cdot L$	$L6a < 0.167 \cdot L$
$L = 105$	$L6 + L5 + L4 > 0.167 \cdot Lb$	$L6 + L5 + L4 > 0.333 \cdot Lb$
$L = 120$	$L2a + L2b < 0.500 \cdot Lb$	$L2a + L2b < 0.500 \cdot Lb$
$L = 135$	$L1a + L1b = 0$	$L1a + L1b = 0$
$L = 150$	$TSgc > 1500$	$TSgc > 1800$
	$LGP_{t=5} = 365$	$LGP_{t=5} = 365$
<b>Barley (sub-tropics)</b>	<i>Climates: STr (WR + SR)</i>	<i>Climates: STr (WR + SR)</i>
(Barley III)	$L6a < 0.333 \cdot L$	$L6a < 0.167 \cdot L$
$L = 90$	$L6 + L5 + L4 > 0.167 \cdot Lb$	$L6 + L5 + L4 > 0.333 \cdot Lb$
$L = 105$	$L2a + L2b < 0.500 \cdot Lb$	$L2a + L2b < 0.500 \cdot Lb$
$L = 120$	$L1a + L1b = 0$	$L1a + L1b = 0$
$L = 135$	$TSgc > 1400$	$TSgc > 1700$
	$LGP_{t=5} = 365$	$LGP_{t=5} = 365$
<b>Wheat (tropics)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
(Wheat IV)	$L6a < 0.333 \cdot L$	$L6a < 0.167 \cdot L$
$L = 100$ at 20.0°C	$L6 + L5 + L4 > 0.167 \cdot Lb$	$L6 + L5 + L4 > 0.333 \cdot Lb$
$L = 130$ at 17.5°C	$L2a + L2b < 0.500 \cdot Lb$	$L2a + L2b < 0.500 \cdot Lb$
$L = 160$ at 15.0°C	$L1a + L1b = 0$	$L1a + L1b = 0$
$L = 190$ at 12.5°C	$TSgc > 1600$	$TSgc > 2000$
	$LGP_{t=5} = 365$	$LGP_{t=5} = 365$
<b>Barley (tropics)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
(Barley IV)	$L6a < 0.333 \cdot L$	$L6a < 0.167 \cdot L$
$L = 100$ at 20.0°C	$L6 + L5 + L4 > 0.167 \cdot Lb$	$L6 + L5 + L4 > 0.333 \cdot Lb$
$L = 130$ at 17.5°C	$L2a + L2b < 0.500 \cdot Lb$	$L2a + L2b < 0.500 \cdot Lb$
$L = 160$ at 15.0°C	$L1a + L1b = 0$	$L1a + L1b = 0$
$L = 190$ at 12.5°C	$TSgc > 1500$	$TSgc > 1900$
	$LGP_{t=5} = 365$	$LGP_{t=5} = 365$

\* applicable to post-dormancy period only

Crop	Sub-optimal Conditions	Optimal Conditions
<b>Indica Rice (wetland)</b>	<i>Climates: Tr, STR (SR+WR)</i>	<i>Climates: Tr, STR (SR+WR)</i>
<i>L = 105/120/135/150</i>	<i>L5=L6=0</i>	<i>L6=L5=L4 = 0</i>
	<i>L3 + L2+ L1 &gt;0.833*L</i>	<i>L3+L2+L1 = 1.000*L</i>
	<i>L4a &lt; 0.167*L</i>	<i>L1 &lt; 0.667*L</i>
	<i>L4b &lt; 0.167*L</i>	<i>TSgc &gt;3000</i>
	<i>TSgc&gt;2400</i>	
<b>Indica Rice (dryland)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 105/120/135</i>	<i>L5=L6=0</i>	<i>L6=L5=L4 = 0</i>
	<i>L3 + L2+ L1 &gt;0.833*L</i>	<i>L3+L2+L1 = 1.000*L</i>
	<i>L4a &lt; 0.167*L</i>	<i>L1 &lt; 0.667*L (not implemented)r</i>
	<i>L4b &lt; 0.167*L</i>	<i>TSgc &gt;3000</i>
	<i>TSgc&gt;2400</i>	
<b>Japonica Rice (wetland)</b>	<i>Climates: Tr, STR (SR+WR), Te</i>	<i>Climates: Tr, STR (SR+WR), Te</i>
<i>L = 105/120/135/150</i>	<i>L5a+L4a &lt; 0.400*L</i>	<i>L5a+L4a &lt; 0.400*L</i>
	<i>L4 &gt;0</i>	
	<i>L2a+L2b &lt; 0.667*L</i>	<i>L2a+L2b &lt; 0.667*L</i>
	<i>L1a+L1b &lt; 0.200*L</i>	<i>L1a+L1b &lt; 0.200*L</i>
	<i>L4b+L5b &lt; 0.250*L</i>	<i>L4b+L5b &lt; 0.250*L</i>
	<i>L6=0</i>	<i>L6=0</i>
	<i>TSgc &gt; 1800</i>	<i>TSgc&gt; 2200</i>
<b>Lowland Maize and Sorghum</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
(Maize I)	<i>L4a+L4b &lt; 0.167*L</i>	<i>L4 = 0</i>
(Sorghum I)	<i>TSgc &gt; 2200</i>	<i>TSgc &gt; 2500</i>
<i>L = 90/105/120/135</i>		
<b>Highland Maize and Sorghum</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
(Maize II)	<i>L1 = 0</i>	<i>L1 = 0</i>
(Sorghum II)	<i>L2 = 0</i>	<i>L2 = 0</i>
<i>L = 105 at 20.0°C</i>	<i>L5 &lt; 0.500*L</i>	<i>L5 &lt; 0.333*L</i>
<i>L = 180 at 17.5°C</i>	<i>L6=0</i>	<i>L6=0</i>
<i>L = 300 at 15.0°C</i>	<i>L3a+L3b &lt; 0.333L</i>	<i>L3a+L3b &lt; 0.333*L</i>
	<i>TSgc &gt; 2200</i>	<i>TSgc &gt; 2500</i>
<b>Maize (sub-tropics, grain)</b>	<i>Climates: STR (SR+WR), Te</i>	<i>Climates: STR (SR+WR), Te</i>
(Maize III)	<i>L5a+L5b &lt; 0.200*L</i>	<i>L5 = &lt;0.200*L</i>
<i>L = 105/120/135/150/165/180</i>	<i>L6=0</i>	<i>L6=0</i>
	<i>TSgc&gt;1900</i>	<i>TSgc&gt;2400</i>
<b>Maize (temperate, silage)</b>	<i>Climates: STR (SR+WR), Te</i>	<i>Climates: STR (SR+WR), Te</i>
(Maize IV)	<i>L5 &lt; 0.667*L</i>	<i>L5 &lt; 0.667*L</i>
<i>L = 105/120/135/150/165/180</i>	<i>L6=0</i>	<i>L6=0</i>
	<i>TSgc&gt;1700</i>	<i>TSgc &gt; 1900</i>
	<i>no permafrost</i>	<i>no permafrost</i>
<b>Irish Potato</b>	<i>Climates: B, Te, STR, Tr</i>	<i>Climates: B, Te, STR, Tr</i>
<i>L = 90/120/150/180</i>	<i>L6a &lt; 0.333*L</i>	<i>L6a &lt; 0.333*L</i>
	<i>L6b &lt; 0.167*L</i>	<i>L6b &lt; 0.167*L</i>
	<i>L2 &lt; 0.333*L</i>	<i>L2 &lt; 0.333*L</i>
	<i>L1 = 0</i>	<i>L1 = 0</i>
	<i>L6+L5+L4 &gt; 0.400*L</i>	<i>L6+L5+L4 &gt; 0.400*L</i>
	<i>TSgc &gt;1200</i>	<i>TSgc&gt;1500</i>
	<i>no permafrost</i>	<i>no permafrost</i>
<b>Pearl Millet</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 70/90</i>	<i>L4a+L4b &lt; 0.167*L</i>	<i>L4 = 0</i>
	<i>L5=L6=0</i>	<i>L5=L6=0</i>
	<i>TSgc &gt; 1600</i>	<i>TSgc &gt; 1800</i>
<b>Cowpea</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 80/100/120</i>	<i>L4a+L4b &lt; 0.167*L</i>	<i>L4 = 0</i>
	<i>L5=L6=0</i>	<i>L5=L6=0</i>
	<i>TSgc &gt; 1800</i>	<i>TSgc &gt; 2100</i>
<b>Cassava</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 12 months</i>	<i>L4 &lt; 0.333*L</i>	<i>L4 &lt; 0.167*L</i>
	<i>L1 &lt; 0.667 * L</i>	<i>L1 &lt; 0.667* L</i>
	<i>L5=L6=0</i>	<i>L5=L6=0</i>
	<i>TSgc&gt; 6500</i>	<i>TSgc&gt; 7500</i>

Crop	Sub-optimal Conditions	Optimal Conditions
<b>Sweet Potato</b>	<i>Climates: Tr, STr (SR)</i>	<i>Climates: Tr, STr (SR)</i>
<i>L = 150/180/210</i>	$L4 < 0.333 * L$	$L4 < 0.333 * L$
	$L5 = L6 = 0$	$L5 = L6 = 0$
	$L1 < 0.500 * L$	$L1 < 0.500 * L$
	$TSgc > 3500$	$TSgc > 4000$
<b>Foxtail Millet (Setaria)</b>	<i>Climates: STr (SR+WR), Te</i>	<i>Climates: STr (SR+WR), Te</i>
<i>L = 75/90/105/120</i>	$L5a + L5b < 0.167 * L$	$L5 = 0$
	$L3 > 0$	$L3 > 0$
	$L6 = 0$	$L6 = 0$
	$TSgc > 1500$	$TSgc > 1800$
<b>Sugarcane</b>	<i>Climates: Tr, STr (SR + WR)</i>	<i>Climates: Tr, STr (SR + WR)</i>
<i>L = 12 months</i>	$L4 + L5 < 0.667 * L$	$L4 < 0.667 * L$
	$L5 + L4 + L3 > 0.084 * L$ (n.a)	$L4 + L3 > 0.084 * L$
	$L6 = 0$	$L5 + L6 = 0$
	$TSgc > 6250$	$TSgc > 6500$
<b>Sugarbeet</b>	<i>Climates: Te, STr (SR+WR)</i>	<i>Climates: Te, STr (SR+WR)</i>
<i>L = 150/165/180/195/210</i>	$L6a < 0.333 * L$ ; $L6b < 0.167 * L$	$L6a < 0.333 * L$ ; $L6b < 0.167 * L$
	$L6 + L5 > 0.167 * L$	$L6 + L5 > 0.167 * L$
	$L2 < 0.400 * L$	$L2 < 0.400 * L$
	$L1 = 0$	$L1 = 0$
	$TSgc > 1600$	$TSgc > 2000$
	no permafrost	no permafrost
<b>Lowland Phaseolus Bean</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 90/120/150</i>	$L4a + L4b < 0.167 * L$	$L4 = 0$
	$L6 + L5 = 0$	$L6 + L5 = 0$
	$TSgc > 1800$	$TSgc > 2400$
<b>Highland Phaseolus Bean</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 120 at 17.5°C</i>	$L5a + L5b < 0.500 * L$	$L5a + L5b < 0.500 * L$
<i>L = 150 at 15.0°C</i>	$L3a + L3b < 0.050 * L$	$L3a + L3b < 0.050 * L$
<i>L = 180 at 12.5°C</i>	$L1 + L2 = 0$	$L1 + L2 = 0$
	$L6 = 0$	$L6 = 0$
	$TSgc > 1500$	$TSgc > 1800$
<b>Phaseolus Bean (temperate)</b>	<i>Climates: Te, STr (SR+WR)</i>	<i>Climates: Te, STr (SR+WR)</i>
<i>L = 90/120/150</i>	$L5 < 0.667 * L$	$L5 < 0.667 * L$
	$L5 + L4 + L3 > 0.500 * L$	$L5 + L4 + L3 > 0.500 * L$
	$L1 = 0$	$L1 = 0$
	$L6 = 0$	$L6 = 0$
	$TSgc > 1400$	$TSgc > 1700$
<b>Chickpea (cold tolerant)</b>	<i>Climates: STr (WR)</i>	<i>Climates: STr (WR)</i>
<i>L = 150/180</i>	$L6a < 0.333 * L$	$L6a < 0.333 * L$
<i>Winter sowing (cold resistant)</i>	$L5b + L6b < 0.167 * L$	$L5b + L6b < 0.167 * L$
<i>(data from ICARDA)</i>	$L6a + L5a + L4a > 0.333 * L$	$L6a + L5a + L4a > 0.333 * L$
	$L4b + L3b + L2b = 0$	$L4b + L3b + L2b = 0$
	$L1 < 0.167 * L$	$L1 < 0.167 * L$
	$L3a > 0$	$L3a > 0$
	$TSgc > 1700$	$TSgc > 2200$
<b>Chickpea</b>	<i>Climates: Tr, STr (SR+WR)</i>	<i>Climates: Tr, STr (SR+WR)</i>
<i>L = 90/105/120</i>	$L6a < 0.167 * L$	$L6 = 0.167$
	$L1 < 0.167 * L$	$L1 < 0.167 * L$
	$L6b + L5b + L4b + L3b + L2b = 0$	$L5b + L4b + L3b + L2b = 0$
	$L6a + L5a + L4a > 0.167 * L$	$L6a + L5a + L4a > 0.167 * L$
	$L3a > 0$	$L3a > 0$
	$TSgc > 1500$	$TSgc > 2100$
<b>Groundnut</b>	<i>Climates: Tr, STr (SR+ WR), Te</i>	<i>Climates: Tr, STr (SR+ WR), Te</i>
<i>L = 90/105/120</i>	$L4a + L4b < 0.167 * L$	$L4 = 0$
	$L5 = L6 = 0$	$L5 = L6 = 0$
	$TSgc > 2200$	$TSgc > 2500$
<b>Soybean (tropics)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 90/105/120</i>	$L4a + L4b < 0.500 * L$	$L4 < 0.333 * L$
	$L5 = L6 = 0$	$L5 = L6 = 0$
	$TSgc > 2100$	$TSgc > 2400$

Crop	Sub-optimal Conditions	Optimal Conditions
<b>Soybean (sub-tropics)</b>	<i>Climates: STr(SR+WR), Te</i>	<i>Climates: STr(SR+WR), Te</i>
<i>L = 105/120/135</i>	<i>L5 = 0</i>	<i>L5 = 0</i>
	<i>L3+L4 &gt; 0.333*L</i>	<i>L6 = 0</i>
	<i>L1 &lt; 0.333*L</i>	<i>L3+L4 &gt; 0.333*L</i>
	<i>TSgc &gt; 1700</i>	<i>L1 &lt; 0.333*L</i>
		<i>TSgc &gt; 2000</i>
<b>Sunflower (tropics)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 135/150/165</i>	<i>L5 + L4 + L3 &gt; 0.500 * L</i>	<i>L5 + L4 + L3 &gt; 0.500 * L</i>
	<i>L3 &lt; 0.500*L</i>	<i>L3 &lt; 0.500*L</i>
	<i>L4 &gt; 0.167 * L</i>	<i>L4 &gt; 0.167 * L</i>
	<i>L1 = L2 = L6 = 0</i>	<i>L1 = L2 = L6 = 0</i>
	<i>TSgc &gt; 2200</i>	<i>TSgc &gt; 2500</i>
<b>Sunflower (sub-tropics)</b>	<i>Climates: STr (SR+WR), Te</i>	<i>Climates: STr (SR+WR), Te</i>
<i>L = 135/150/165</i>	<i>L5 &lt; 0.300*L</i>	<i>L5 &lt; 0.200*L</i>
	<i>L5+L4+L3 &gt; 0.400*L (n.a.)</i>	<i>L5+L4+L3 &gt; 0.400*L (n.a.)</i>
	<i>L1 &lt; 0.333*L</i>	<i>L1 &lt; 0.333*L</i>
	<i>L6 = 0</i>	<i>L6 = 0</i>
	<i>TSgc &gt; 2000</i>	<i>TSgc &gt; 2400</i>
<b>Winter Rape</b>	<i>Climates: Te, STr (SR+TR)</i>	<i>Climates: Te, STr (SR+TR)</i>
<i>L = 35 + 105</i>	<i>L6a &lt; 0.667*Lb</i>	<i>L6a &lt; 0.500*Lb</i>
<i>L = 45 + 120</i>	<i>L2a+L2b &lt; 0.400*Lb</i>	<i>L2a+L2b &lt; 0.400*Lb</i>
	<i>L1a+L1b = 0</i>	<i>L1a+L1b = 0</i>
	<i>L2b+L3b+L4b+L5b &lt; 0.500*Lb</i>	<i>L2b+L3b+L4b+L5b &lt; 0.500*Lb</i>
	<i>N4b+N5b+N6b &gt; La</i>	<i>N4b+N5b+N6b &gt; La</i>
	<i>TSgc &gt; 1200 *</i>	<i>TSgc &gt; 1400 *</i>
	<i>LGP t=5 &lt; 365</i>	<i>LGP t=5 &lt; 365</i>
	<i>dormancy required</i>	<i>dormancy required</i>
	<i>no permafrost</i>	<i>no permafrost</i>
<b>Spring Rape</b>	<i>Climates: Te, STr (SR+WR)</i>	<i>Climates: Te, STr (SR+WR)</i>
<i>L = 135/150/165</i>	<i>L6a &lt; 0.333*L</i>	<i>L6a &lt; 0.333*L</i>
	<i>L2a+L2b &lt; 0.400*L</i>	<i>L2a+L2b &lt; 0.400*L</i>
	<i>L1a+L1b = 0</i>	<i>L1a+L1b = 0</i>
	<i>L2b+L3b+L4b+L5b &lt; 0.500*L</i>	<i>L2b+L3b+L4b+L5b &lt; 0.500*L</i>
	<i>TSgc &gt; 1400</i>	<i>TSgc &gt; 1800</i>
	<i>LGP t=5 &lt; 365</i>	<i>LGP t=5 &lt; 365</i>
	<i>no permafrost</i>	<i>no permafrost</i>
<b>Rape (rabi)</b>	<i>Climates: Tr, STr (SR + WR)</i>	<i>Climates: Tr, STr (SR + WR)</i>
<i>L = 135/150/165</i>	<i>L6a &lt; 0.333*L</i>	<i>L6a &lt; 0.333*L</i>
	<i>L2a+L2b &lt; 0.500*L</i>	<i>L2a+L2b &lt; 0.500*L</i>
	<i>L1a+L1b = 0</i>	<i>L1a+L1b = 0</i>
	<i>L6+L5+L4 &gt; 0.167*L</i>	<i>L6+L5+L4 &gt; 0.167*L</i>
	<i>TSgc &gt; 1400</i>	<i>TSgc &gt; 1800</i>
	<i>LGP t=5 = 365</i>	<i>LGP t=5 = 365</i>
<b>Cotton (tropics)</b>	<i>Climates: Tr</i>	<i>Climates: Tr</i>
<i>L = 135/150/165/180</i>	<i>L4a+L4b &lt; 0.250*L</i>	<i>L4 = 0</i>
	<i>TSgc &gt; 3000</i>	<i>TSgc &gt; 3400</i>
<b>Cotton (sub-tropics)</b>	<i>Climates: STr (SR+WR), Te</i>	<i>Climates: STr (SR+WR), Te</i>
<i>L = 135/150/165</i>	<i>L4a+L4b &lt; 0.333*L</i>	<i>L4a+L4b &lt; 0.167*L</i>
	<i>TSgc &gt; 2700</i>	<i>TSgc &gt; 3000</i>
<b>Banana/Plantain</b>	<i>Climate: Tr, STr (SR+WR)</i>	<i>Climate: Tr, STr (SR+WR)</i>
<i>L = 12 months</i>	<i>L5 &lt; 0.667*L</i>	<i>L5 &lt; 0.500*L</i>
	<i>L2+L3 &gt; 0.333*L</i>	<i>L2+L3 &gt; 0.400*L</i>
	<i>L1 &lt; 0.667 * L</i>	<i>L1 &lt; 0.500 * L</i>
	<i>TSgc &gt; 6000</i>	<i>TSgc &gt; 7000</i>
<b>Oil Palm</b>	<i>Climate: Tr, STr (SR)</i>	<i>Climate: Tr, STr (SR)</i>
<i>L = 12 months</i>	<i>L4a+L4b &lt; 0.333*L</i>	<i>L4 &lt; 0.167*L</i>
	<i>TSgc &gt; 7500</i>	<i>TSgc &gt; 8000</i>

\* applicable to post-dormancy period only

Crop	Sub-optimal Conditions	Optimal Conditions
<b>Olive</b>	<i>Climates: STR (SR+WR), Te</i>	<i>Climates: STR (SR+WR), Te</i>
<i>L = 12 months</i>	<i>L8 = L9 = 0</i>	<i>L8 = L9 = 0</i>
	<i>L7 + L6 + L5 + L4 &gt; 0.400*L</i>	<i>L7 + L6 + L5 + L4 &gt; 0.400*L</i>
	<i>L4 + L3+ L2+ L1 &gt; 0.333*L</i>	<i>L4 + L3+ L2+ L1 &gt; 0.333*L</i>
	<i>TSgc &gt; 4000</i>	<i>TSgc &gt; 5000</i>
<b>Alfalfa (Lucerne)</b>	<i>Climates: Te, STR (SR+WR)</i>	<i>Climates: Te, STR (SR+WR)</i>
<i>L = LGP<sub>t=5</sub></i>	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt; 120 days .</i>	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt; 150 days .</i>
	<i>L6 &lt; 0.500*L</i>	<i>L6 &lt; 0.500*L</i>
	<i>L1 &lt; 0.333*L</i>	<i>L1 &lt; 0.333*L</i>
	<i>TSgc &gt; 1800</i>	<i>TSgc &gt; 2250</i>
<b>Grasses</b>	<i>Climates: Tr, STR (SR+WR), Te</i>	<i>Climates: Tr, STR (SR+WR), Te</i>
<i>L = LGP<sub>t=5</sub></i>	<i>TSgc &gt; 750</i>	<i>TSgc &gt; 1000</i>
	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt;90 days</i>	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt;120 days</i>
<b>Grasses</b>	<i>Climates: Te, B</i>	<i>Climates: Te</i>
<i>L = LGP<sub>t=5</sub></i>	<i>TSgc &gt; 750</i>	<i>TSgc &gt; 1000</i>
	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt;90 days</i>	<i>Gc (L) = (LGP<sub>t=5</sub>) &gt;120 days</i>